

Product Data Sheet

Purified anti-human CD178 (Fas-L)

Catalog # / Size: 306401 / 25 µg
306402 / 100 µg

Clone: NOK-1

Isotype: Mouse IgG1, κ

Workshop Number: VII 70322

Immunogen: L5178Y mouse T lymphoma cells expressing recombinant human FasL


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human Fas Ligand transfected cells stained with purified NOK-1, followed by biotinylated anti-mouse IgG and Sav-PE

Applications:

Applications: FC - Quality tested
IP, IF, ICC, WB - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10^6 cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,2}, immunofluorescence microscopy³, immunocytochemistry², blocking of Fas induced apoptosis¹, and Western blotting¹¹. Fas Ligand is expressed at low density on activated cells. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 306407) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated antibody (Cat. No. 306404) or biotinylated anti-mouse IgG second step (Cat. No. 405303), followed by SAV-PE (Cat. No. 405204)). In addition, applying matrix metalloproteinases (MMPs) inhibitor in the cell culture system will increase the FasL staining intensity. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 306409).

Application References:

- Kayagaki N, *et al.* 1995. *J. Exp. Med.* 182:1777.
- Andreola G, *et al.* 2002. *J. Exp. Med.* 195:1303.
- Bossi G, *et al.* 1999. *Nature Medicine* 5:90.
- Strauss L, *et al.* 2009. *J. Immunol.* 182:1469. PubMed
- Li JH, *et al.* 2009. *Am J. Pathol.* 175:1124. PubMed
- Kruger K, *et al.* 2011. *J. Appl. Physiol.* 110:1226. PubMed
- Qin G, *et al.* 2012. *J. Infect. Dis.* PubMed
- Khalid M, *et al.* 2012. *J. Virol.* 86:4906. PubMed
- Qin G, *et al.* 2012. *J. Infect. Dis.* 205:1646. PubMed
- Shrestha B, *et al.* 2012. *J. Virol.* 86:8937. PubMed
- Mooren FC, *et al.* 2012. *J. Appl. Physiol.* 113:1082. PubMed
- Herr I, *et al.* 2000. *Cell Death Differ.* 7:129. (WB)

Description: CD178 is a 38-42 kD type II glycoprotein also known as Fas ligand and CD95L. CD178 belongs to the TNF superfamily and is expressed on activated T lymphocytes, NK cells, monocytes, and granulocytes. CD178 is also expressed on parenchymal cells of the retina and cornea, retinal pigment epithelial cells, and testis. The extracellular region of FasL can be cleaved by matrix metalloproteinases (MMPs) to give rise to a 26 kD soluble protein. CD178 binds to CD95, a member of the TNFR superfamily, to induce apoptosis. CD95/CD95L interactions play an important role in the maintenance of peripheral tolerance and survival.

Antigen References:

- Suda T, *et al.* 1997. *J. Exp. Med.* 12:204.
- Kayagaki N, *et al.* 1995. *J. Exp. Med.* 182:1777.
- Tanaka M, *et al.* 1995. *EMBO J.* 14:1129.

Related Products:

Product
Purified anti-human CD95 (FAS)
Purified anti-human CD95 (FAS)
Purified Mouse IgG1, κ Isotype Ctrl
APC Goat anti-mouse IgG (minimal x-reactivity)

Clone
DX2
EOS9.1
MOPC-21
Poly4053

Application
FC, IF, IHC
FC
FC, ICFC, ICC, IF, IHC, IP, WB
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

HRP Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	ELISA, IHC, WB
PE Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Purified anti-Cytochrome c	2.7D5	FC
Purified anti-PARP	5A5	IF, WB
Purified anti-Cytochrome c	6H2.B4	IP, ICFC, IF, ICC
Purified anti-Cytochrome c	7H8.2C12	WB
Purified anti-Caspase-8	4-1-20	WB



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 9727 Pacific Heights Blvd, San Diego, CA 92121 www.biollegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587