

## ChargeSwitch®-Pro Filter Plasmid Miniprep Kit

Cat. no. CS31102

Size: 10 Miniprep reactions

Store at room temperature

### Description

The ChargeSwitch®-Pro Filter Plasmid Miniprep Kit contains all the components required for the rapid and efficient isolation of highly pure plasmid DNA from *E. coli* cells. The purification columns in the kit contain a novel ChargeSwitch®-derivatized membrane that is positively charged at low pH and neutral at pH 8.5, to bind and elute plasmid DNA without the use of harsh reagents.

Using the kit, you prepare cell lysates with a modified alkaline lysis procedure and then purify the plasmid DNA using a simple centrifugation- or vacuum-based protocol. In low pH conditions, the ChargeSwitch®-derivatized membrane binds the negatively charged nucleic acid backbone. Proteins and other contaminants are not bound and simply wash away in the aqueous wash buffers.

To elute the DNA, the charge of the membrane is neutralized by raising the pH to 8.5 using a low-salt elution buffer. The purified plasmid DNA is ready for use in your downstream application of choice, including mammalian transfection, automated fluorescent DNA or manual sequencing, PCR, cloning, *in vitro* transcription, bacterial cell transformations, or restriction digestion.

The ChargeSwitch®-Pro Filter Plasmid Miniprep Kit offers the following advantages:

- High-quality, high-yield plasmid DNA purification from *E. coli* without the use of ethanol, chaotropic salts, or organic solvents.
- Designed to isolate plasmid DNA from samples using a simple centrifugation or vacuum protocol following sample preparation.
- Reliable performance of the purified plasmid DNA in a variety of applications, including mammalian cell transfection, automated and manual sequencing, amplification reactions, *in vitro* transcription, bacterial cell transformation, cloning, and labeling.
- Compatible for use in isolation of plasmid from *endA*+ strains.


### Kit Contents

Component	Quantity
ChargeSwitch®-Pro Resuspension Buffer (10 mM Tris-HCl, pH 8.5, 10 mM EDTA)	4 ml
RNase A	80 µl
ChargeSwitch®-Pro Lysis Buffer	4 ml
ChargeSwitch®-Pro Precipitation Buffer	4 ml
ChargeSwitch®-Pro Wash Buffer 1	10 ml
ChargeSwitch®-Pro Wash Buffer 2	4 ml
ChargeSwitch®-Pro Elution Buffer (10 mM Tris-HCl, pH 8.5)	4 ml
ChargeSwitch®-Pro Filter Plasmid Miniprep Columns	10
ChargeSwitch®-Pro Elution Tubes	10

### Specifications

The ChargeSwitch®-Pro Filter Plasmid Miniprep Kit employs a nested column design consisting of an inner flat bottomed column (A), which fits into an outer luer bottomed column (B). The column assembly is provided in a 2 ml Collection Tube. The inner column (Lysate Clarification Column) is used for rapid clarification of the bacterial lysate. The outer column (Binding Column) contains the ChargeSwitch® derivitized membrane which binds plasmid DNA from the clarified lysate.

	Miniprep
Starting Material (fresh, overnight LB culture)	5 ml
Binding Capacity (per column)	40 µg
Recommended Elution Volume	50–150 µl
Typical DNA Yield	25 µg
Plasmid Size	3–9 kb
Purity OD 260/280	>1.8
Purity OD 260/230	>1.8
Endotoxin	≤2 EU/µg DNA



### Shipping and Storage

All components are shipped and stored at room temperature. **Do not freeze the columns.**

## Important Guidelines and Parameters

### Bacterial Cultures

- Grow transformed *E. coli* in LB medium with the appropriate antibiotic.
- **Do not** use rich medium (e.g., Terrific Broth) to grow the *E. coli*.
- Use overnight bacterial cultures with an absorbance of 1–2 OD at 600 nm ( $A_{600}$ ).
- For best results, use fresh overnight cultures. However, the kit can also be used to purify DNA from frozen cell pellets.
- Use up to 5 ml of bacterial culture for minipreps.

### Handling DNA

- Maintain a sterile environment and use sterile equipment when handling DNA to avoid DNase contamination.
- Ensure that no DNase is introduced into the solutions supplied with the kit.

### Columns

- **Do not freeze the columns.** Freezing may damage the ChargeSwitch®-derivatized membrane.

### Before Starting

- For a new kit, add the entire volume of the provided RNase A to the ChargeSwitch®-Pro Resuspension Buffer and mix.
- If necessary, warm the Lysis Buffer to 37°C to dissolve any precipitate.
- If the ambient temperature is >25°C, chill the ChargeSwitch®-Pro Precipitation Buffer on ice before use to improve results.
- For best results, use the Elution Buffer provided in the kit. **Do not elute in water.** If you need to elute in any other buffer, be sure to use a buffer of **pH 8.5–9.0**. If the pH of the buffer is <8.5, the DNA will not elute efficiently.

### Materials Needed

- Overnight bacterial culture (see **Bacterial Cultures**, above)
- Microcentrifuge
- Vacuum protocol only: Vacuum manifold and pump (capable of producing pressure of 15–20 in. Hg or –500 to –700 mbar). Invitrogen produces the EveryPrep™ Universal Vacuum Manifold (Invitrogen Cat. no. K211101), which is ideal for this application.
- Adjustable pipettes and aerosol barrier pipette tips

### Preparing the Sample

1. Harvest up to 5 ml of overnight bacterial culture by centrifugation at  $\geq 8,500 \times g$  for 10 minutes.
2. Resuspend the cell pellet in 250  $\mu$ l of ChargeSwitch®-Pro Resuspension Buffer premixed with RNase A (see **Before Starting**, above). Invert the tube or vortex until all cell clumps are dispersed.
3. Add 250  $\mu$ l ChargeSwitch®-Pro Lysis Buffer. Mix by inverting the capped tube 5–10 times until the solution becomes homogenous. **Do not vortex.**
4. Incubate at room temperature for 5 minutes. The lysate turns clear and viscous. **Do not incubate longer than 5 minutes.**
5. Add 250  $\mu$ l ChargeSwitch®-Pro Precipitation Buffer. Mix by inversion 6–10 times until the precipitate that forms becomes homogeneous. **Do not vortex.**
6. Proceed to **Centrifugation Protocol**, **Vacuum Protocol**, or **EveryPrep™ Universal Vacuum Manifold Protocol**.

### Centrifugation Protocol

Follow the steps below to purify plasmid DNA from overnight cultures using a swinging bucket centrifuge. All steps are performed at room temperature.

#### Binding the DNA

1. Transfer the lysate (**Preparing the Sample**, Step 5) onto the ChargeSwitch®-Pro Filter Mini Column.
2. Incubate for 1 minute at room temperature to allow precipitate to float to the surface.
3. Centrifuge the column at  $1,350 \times g$  for 1 minute. Remove the column assembly and decant the flow-through.  
**Note:** If the lysate has not completely passed through the column after the initial spin, repeat centrifugation for 1 minute at  $8,500 \times g$ .
4. Remove and discard the Lysate Clarification Column (see page 1). Re-insert the Binding Column into the tube.

#### Washing the Column

1. Add 750  $\mu$ l of ChargeSwitch®-Pro Wash Buffer 1 to the column.
2. Centrifuge at  $8,500 \times g$  for 1 minute. Remove the column and discard the flow-through. Re-insert the column into the tube.
3. Add 250  $\mu$ l of ChargeSwitch®-Pro Wash Buffer 2 to the column.
4. Centrifuge at  $8,500 \times g$  for 1 minute. Remove the column. Discard the flow-through *and* the tube.
5. Insert the column into a clean 1.7 ml Elution Tube, and proceed to **Eluting the DNA**, next page.

## Vacuum Protocol

Follow the steps below to purify plasmid DNA from overnight cultures using a luer lock vacuum manifold, pump, and microcentrifuge. All steps are performed at room temperature. For a vacuum assisted elution protocol, see the **EveryPrep™ Universal Vacuum Manifold Protocol** below.

### Binding the DNA

1. Discard the 2 ml Collection Tube from the ChargeSwitch®-Pro Filter Mini Column assembly.
2. Insert the ChargeSwitch®-Pro Filter Mini Column into the luer extension of a vacuum manifold.
3. Transfer the lysate (**Preparing the Sample**, Step 5) onto the column.
4. Incubate for 1 minute at room temperature to allow precipitate to float to the surface.
5. Apply maximum vacuum pressure (15–20 in. Hg) until the liquid has passed through the column.
6. Release the vacuum. Gently remove the inner Lysate Clarification Column (see page 1) and discard.

### Washing the Column

1. Add 750 µl of ChargeSwitch®-Pro Wash Buffer 1 to the column.
2. Apply maximum vacuum pressure until the liquid has passed through the column. Release the vacuum.
3. Add 250 µl of ChargeSwitch®-Pro Wash Buffer 2 to the column.
4. Apply maximum vacuum pressure until the liquid has passed through the column.
5. Remove the column from the manifold and insert it into a clean 1.7 ml Elution Tube. Proceed to **Eluting the DNA**, below.

### Eluting the DNA

1. Add 50–150 µl of ChargeSwitch®-Pro Elution Buffer onto the column and incubate for 1 minute.
2. Centrifuge at  $8,500 \times g$  for 1 minute. Discard the Binding Column. **Do not** reuse.
3. The eluate contains the purified plasmid DNA. Store plasmid DNA at 4° C for immediate use or at –20° C for long-term storage. Avoid repeated freeze-thawing of purified DNA. Calculate DNA yield by UV absorbance at 260 nm.

## EveryPrep™ Universal Vacuum Manifold Protocol

Follow the steps below to purify plasmid DNA from overnight cultures using the EveryPrep™ Universal Vacuum Manifold and vacuum pump. All steps are performed at room temperature. For alternative protocols using the EveryPrep™ Universal Vacuum Manifold, refer to the manual.

### Binding the DNA

1. Discard the 2 ml Collection Tube from the ChargeSwitch®-Pro Filter Mini Column assembly.
2. Place the Waste Tray in one chamber of the manifold.
3. Seat the Mini Elution Top Plate above the Waste Tray and insert the required number of assembled columns firmly into the plate. Block the remaining holes with the provided stoppers.
4. Transfer the lysate (**Preparing the Sample**, Step 5) onto the column.
5. Incubate for 1 minutes at room temperature to allow precipitate to float to the surface.
6. Apply maximum vacuum pressure (15–20 in. Hg) until the liquid has passed through the column.
7. Release the vacuum. Gently remove the inner Lysate Clarification Column (see page 1) and discard.

### Washing the Column

1. Add 750 µl of ChargeSwitch®-Pro Wash Buffer 1 to the column.
2. Apply maximum vacuum pressure until the liquid has passed through the column. Release the vacuum.
3. Add 250 µl of ChargeSwitch®-Pro Wash Buffer 2 to the column.
4. Apply maximum vacuum pressure until the liquid has passed through the column. Release the vacuum.

### Eluting the DNA by Vacuum

1. Place the Elution Rack into the Elution Chamber. The Elution Rack should contain a clean 1.7 ml Elution Tube in each position corresponding to a ChargeSwitch®-Pro Filter Mini Column.
2. Transfer the Mini Elution Top Plate so that it is positioned over the Elution Chamber.
3. Add 75–150 µl of ChargeSwitch®-Pro Elution Buffer onto each column and incubate for 1 minute.
4. Apply a vacuum of 5–10 in. Hg for 2 minutes to allow the elution buffer to completely pass through the column before releasing the vacuum.
5. Remove the Mini Elution Top Plate, and discard the columns.
6. The eluate contains the purified plasmid DNA. Store purified plasmid DNA at 4° C for immediate use or at –20° C for long-term storage.

## Troubleshooting

Problem	Possible Cause	Solution
Low plasmid DNA yield	Poor quality of starting material or incomplete lysis	<ul style="list-style-type: none"> <li>• Ensure media is completely removed after cell harvest.</li> <li>• If the cell lysate is too viscous, reduce the amount of cells used per sample. Attempting to lyse too many cells may result in incomplete lysis. See the culture volume recommendations on page 2.</li> <li>• Check the growth conditions of the cell culture to ensure plasmid propagation. Use a high copy number plasmid, if possible.</li> <li>• Cell cultures that are overgrown (<i>e.g.</i>, grown &gt;16 hours) may begin to lyse, resulting in reduced yields and contaminating genomic DNA</li> <li>• Ensure complete resuspension of the bacterial cell pellet. Decrease the amount of starting material used.</li> <li>• Chill the Precipitation Buffer to 4°C before use to improve the precipitation efficiency and plasmid DNA yield.</li> <li>• Mix lysate thoroughly (&gt;10 inversions) to ensure complete lysis.</li> <li>• Increase the incubation time during lysis but do not exceed 5 minutes.</li> </ul>
	Elution conditions require optimization	If you are using a different buffer for elution, ensure that the pH of the buffer is 8.5-9.0.
	ChargeSwitch®-derivatized membrane is not functional	<b>Do not freeze the columns.</b> Store the columns at room temperature. Do not re-use the columns.
	Cell cultures are overgrown	Cells grown more than 16 hours may begin to lyse, resulting in reduced yields
	Quantitation is inaccurate: Supercoiled plasmid DNA fluoresces less than the DNA standard provided in quantitation kit	We recommend using plasmid DNA as a standard when calculating yield using a fluorescence-based DNA quantitation kit. Such kits typically provide non-supercoiled DNA as a standard, which fluoresces more brightly than supercoiled plasmid DNA, leading to inaccuracies in quantitation.
Lysate Clarification Column clogging	Too much precipitate	<ul style="list-style-type: none"> <li>• Incubate lysate for 2–3 minutes to allow precipitate to float.</li> <li>• Cell culture may be overgrown. See page 2 for details on culturing cells.</li> </ul>
Genomic DNA contamination	Genomic DNA sheared during handling	Gently invert tubes to mix after adding buffers. <b>Do not vortex</b> as it can shear the genomic DNA. To efficiently precipitate the genomic DNA away from the plasmid DNA, the genomic DNA must be intact.
RNA contamination	Insufficient RNase treatment	<ul style="list-style-type: none"> <li>• Ensure RNase A is added to the resuspension buffer.</li> <li>• Add additional RNase A to 100 µl/ml if buffer was stored for &gt;6 months.</li> <li>• Mix sufficiently after addition of lysis buffer for adequate RNase A digestion.</li> </ul>
Plasmid DNA degradation	Incorrect lysis procedure	Incubate the lysate at room temperature for no longer than 5 minutes, because it might begin to denature the DNA.

## Additional Products

<u>Product</u>	<u>Amount</u>	<u>Catalog No.</u>
EveryPrep™ Universal Vacuum Manifold	1 unit	K211101
ChargeSwitch®-Pro Filter Collection Tubes	25 tubes	CS32000
ChargeSwitch®-Pro Filter Plasmid Midiprep Kit	25 reactions	CS31104
ChargeSwitch®-Pro Filter Plasmid Maxiprep Kit	10 reactions	CS31106
	25 reactions	CS31107
ChargeSwitch®-Pro Filter Plasmid Miniprep Kit	10 reactions	CS31102
	100 reactions	CS31103

## Purchaser Notification

### Limited Use Label License No. 5: Invitrogen Technology

The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) not to transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. Invitrogen Corporation will not assert a claim against the buyer of infringement of patents owned or controlled by Invitrogen Corporation which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Invitrogen is willing to accept return of the product with a full refund. For information on purchasing a license to this product for purposes other than research, contact Licensing Department, Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, California 92008. Phone (760) 603-7200. Fax (760) 602-6500. Email: [outlicensing@invitrogen.com](mailto:outlicensing@invitrogen.com).