

Human CCL8 (MCP-2) Recombinant Protein Carrier-Free

Catalog Number: 34-8481

Also known as: C-C Motif Chemokine 8, Monocyte Chemoattractant Protein 2

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Contents: Human CCL8 (MCP-2)
Recombinant Protein Carrier-Free
Catalog Number: 34-8481
Concentration: 0.5 mg/mL
Handling Conditions: For best recovery,
quick-spin vial prior to opening. Use in a
sterile environment.
Source: E. coli expressed amino acids
Gln24-Pro99, accession number NM_005623
Molecular Mass: 8.9 kDa
Purity: > 97%, as determined by SDS-PAGE.
Endotoxin: Less than 0.01 ng/ug cytokine,
as determined by the LAL assay.
Bioactivity: The bioactivity of this protein
was determined by transmigration assay of
human monocytes, with maximum
chemotaxis observed at 100-150 ng/mL.



Formulation: Sterile liquid; 0.1 M glycine, pH 3.0

Temperature Limitation: Store at less than or
equal to -70°C.

Batch Code: Refer to vial

Use By: Refer to vial

Description

CCL8, also known as MCP-2 (Monocyte Chemotactic Protein 2), is a member of the CC- subfamily of chemokines and is most closely related to CCL2 (MCP-1) and CCL7 (MCP-3). All three MCPs are secreted by a variety of cell types in response to inflammatory stimuli and play critical roles in the recruitment of leukocytes to areas of inflammation. While they are potent chemoattractants of monocytes and T cells, CCL7 appears to have the broadest range of activity, as it has also been demonstrated to attract activated NK cells, eosinophils, basophils, and neutrophils. CCL8 has also been demonstrated to attract NK cells, eosinophils, and basophils, but requires higher concentrations to do so. CCL8 signals via the G protein-coupled receptor CCR5, which is shared with other CC-chemokines. CCR5 is the primary co-receptor for HIV entry, which the virus binds through the gp120 envelope protein. All CCR5 ligands demonstrate potent inhibition of virus entry into the cell, both through steric hindrance of gp120-CCR5 interaction, and ligand-mediated receptor internalization.

Applications Reported

Human CCL8 Recombinant Protein Carrier-Free is biologically active.

Applications Tested

The bioactivity of this protein was determined by transmigration assay of human monocytes, with maximum chemotaxis observed at 100-150 ng/mL. The ED50 for this effect is less than or equal to 50 ng/mL, which corresponds to a specific activity of greater than or equal to 2 x 10⁴ Units/mg.

References

Thompson WL, Van Eldik LJ. Inflammatory cytokines stimulate the chemokines CCL2/MCP-1 and CCL7/MCP-3 through NFkB and MAPK dependent pathways in rat astrocytes. Brain Res. 2009 Sep 1;1287:47-57.

Jia T, Serbina NV, Brandl K, Zhong MX, Leiner IM, Charo IF, Pamer EG. Additive roles for MCP-1 and MCP-3 in CCR2-mediated recruitment of inflammatory monocytes during Listeria monocytogenes infection. J Immunol. 2008 May 15; 180(10):6846-53.

Proost P, Wuyts A, Van Damme J. Human monocyte chemotactic proteins-2 and -3: structural and functional comparison with MCP-1. J Leukoc Biol. 1996 Jan;59(1):67-74.

Related Products

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Human CCL8 (MCP-2) Recombinant Protein Carrier-Free

Catalog Number: 34-8481

Also known as: C-C Motif Chemokine 8, Monocyte Chemoattractant Protein 2

RUO: For Research Use Only. Not for use in diagnostic procedures.

14-8198 Human CCL7 (MCP-3) Recombinant Protein

14-8398 Human CCL2 (MCP-1) Recombinant Protein