

An Affymetrix Company

# **IHC Antigen Retrieval Solution – High pH (10X)**

Catalog Number: 00-4956

**GPR: General Purpose Reagents. For Laboratory Use.** 

#### **Product Information**

Contents: IHC Antigen Retrieval Solution -

High pH (10X)

REF Catalog Number: 00-4956

Formulation: 10X: 100 mM Tris, 10 mM EDTA

solution, pH 9.0

Temperature Limitation: Store at room

temperature.

Batch Code: Refer to vial
Use By: Refer to vial



The IHC Antigen Retrieval Solution- High pH (10X) is designed for use during the heat induced epitope retrieval (HIER) step prior to immunohistochemistry on formalin-fixed paraffin embedded tissue sections. The use of this buffer in combination with heat (often by microwave, water bath, or pressure cooker) has been shown to restore the antigenicity of proteins modified during the formalin fixation of tissue. This buffer is supplied as a 10X stock solution. Dilution of this buffer to 1X using reagent grade water results in a 10 mM Tris, 1 mM EDTA solution, pH 9.0.

#### **Applications Reported**

IHC Antigen Retrieval Solution – High pH (10X) has been reported for use in immunohistochemical staining of formalin-fixed paraffin embedded tissue sections.

### **Applications Tested**

IHC Antigen Retrieval Solution – High pH (10X) has been tested immunohistochemistry on formalin-fixed paraffin embedded tissue.

## References

Lyck L, Dalmau I, Chemnitz J, Finsen B, Schrøder HD. Immunohistochemical markers for quantitative studies of neurons and glia in human neocortex. J Histochem Cytochem. 2008 Mar;56(3):201-21

Shi SR, Cote RJ, Taylor CR. Antigen retrieval techniques: current perspectives. J Histochem Cytochem. 2001 Aug;49(8):931-7.

Shi SR, Key ME, Kalra KL. Antigen retrieval in formalin-fixed, paraffin-embedded tissues: an enhancement method for immunohistochemical staining based on microwave oven heating of tissue sections. J Histochem Cytochem. 1991 Jun;39(6):741-8

#### **Related Products**

00-4955 IHC Antigen Retrieval Solution - Low pH (10X)