

Product Data Sheet

LEAF™ Purified anti-human HLA-A,B,C

Catalog # / Size: 311411 / 50 µg
 311412 / 500 µg
 311423 / 1 mg

Clone: W6/32

Isotype: Mouse IgG2a, κ


Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Cattle (Bovine, Cow), Cat (Feline)

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.



Human peripheral blood lymphocytes stained with LEAF™ purified W6/32, followed by anti-mouse IgG FITC

Applications:

Applications: FC - *Quality tested*
 IP, WB, IHC, Block, Activ - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 2.0 µg per 10⁶ cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone W6/32 recognizes a monomorphic epitope on the 45 kD polypeptide products of HLA-A, B, C¹⁸.

Additional reported applications (for the relevant formats) include: immunoprecipitation², Western blotting (non-reducing)³, immunohistochemical staining of acetone-fixed frozen tissue sections^{4,5}, blocking^{6,7}, inhibition of NK cell-mediated lysis¹⁰ and activation^{8,9}. Clone W6/32 has been reported not to be suitable for immunohistochemistry on paraffin sections¹⁷. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 311412). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 311428) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

- Application References:**
1. Darrow TL, et al. 1989. *J. Immunol.* 142:3329.
 2. Stern P, et al. 1987. *J. Immunol.* 138:1088.
 3. Tran TM, et al. 2001. *Immunogenetics* 53:440.
 4. Barbatis C, et al. 1981. *Gut* 22:985.
 5. Ayyoub M, et al. 2004. *Cancer Immunity* 4:7.
 6. DeFelice M, et al. 1990. *Cell. Immunol.* 126:420.
 7. Fayen J, et al. 1998. *Int. Immunol.* 10:1347.
 8. Turco MC, et al. 1988. *J. Immunol.* 141:2275.
 9. Geppert TD, et al. 1989. *J. Immunol.* 142:3763.
 10. Wooden SL, et al. 2005. *J. Immunol.* 175:1383.
 11. Nagano M, et al. 2007. *Blood* 110:151.
 12. McLoughlin RM, et al. 2008. *J. Immunol.* 181:1323. PubMed
 13. Takahara M, et al. 2008. *J. Leukoc. Biol.* 83:742. PubMed
 14. Lunemann A, et al. 2008. *J. Immunol.* 181:6170. PubMed
 15. Laing BJ, et al. 2010. *J. Thorac Cardiovasc Surg.* 139:1402. PubMed
 16. Yoshino N, et al. 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 17. Vambutas A, et al. 2000. *Clin. Diagn. Lab. Immun.* 7:79.
 18. Coppieters KT, et al. 2012. *J. Exp. Med.* 209:51. (epitope)
 19. Everds N, et al. 2013. *Toxicol Pathol.* PubMed.

Description: MHC class I antigens associated with β2-microglobulin are expressed by all human nucleated cells. MHC class I molecules are involved in presentation of antigens to CD8⁺ T cells. They play an important role in cell-mediated immune responses and tumor surveillance.

Antigen References: 1. Barclay AN, et al. Eds. 1993. *The Leukocyte Antigen FactsBook*. Academic Press Inc. San Diego.

Related Products:

LEAF™ Purified Mouse IgG2a, κ Isotype Ctrl

Clone

MOPC-173

Application

FC, WB, IP, ICFC, ICC, IF, IHC, FA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.