

Product Data Sheet

Purified anti-human CD257 (BAFF, BLYS)

Catalog # / Size: 318602 / 100 µg

Clone: T7-241

Isotype: Mouse IgG1, κ

Immunogen: recombinant human BAFF(TL7) protein

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

Applications:

Applications: FC - *Quality tested*
ELISA - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: The T7-241 antibody recognizes human BAFF also known as B cell activating factor, tumor necrosis factor ligand superfamily 13B, B lymphocyte stimulator (BLYS), TNF and APOL-related leukocyte expressed ligand 1 (TALL1), and TNF homolog that activates apoptosis, NF-κB, and JNK (THANK). BAFF is a type II transmembrane protein, and a member of the tumor necrosis factor ligand superfamily. BAFF can be proteolytically cleaved to form a soluble protein. BAFF, like other TNF ligand family members, forms homotrimers. The predicted molecular weight of a BAFF monomer is approximately 31 kD. BAFF is expressed on peripheral blood B and T lymphocytes, monocytes, macrophages, and dendritic cells and is upregulated by IFN-γ and downregulated by PMA/ionomycin treatment. The T7-241 antibody has been shown to be useful for flow cytometry and ELISA. BAFF stimulates B and T cell immunity and regulates humoral immunity by binding TACI and BCMA receptors; BAFF prevents apoptosis and promotes B cell survival by binding the BAFF-specific receptor (BAFFR/BR3). The T7-241 antibody has been shown to be useful for flow cytometry and ELISA.

Antigen References: 1. Schneider P, *et al.* 1999. *J. Exp. Med.* 189:1747.
2. Moore PA, *et al.* 1999. *Science* 285:260.
3. Shu HB, *et al.* 1999. *J. Leuko. Biol.* 65:680.
4. Yu G, *et al.* 2000. *Nat. Immunol.* 1:252.
5. Oren DA, *et al.* 2002. *Nat. Struct. Biol.* 9:288.

Related Products:

Product
Purified Mouse IgG1, κ Isotype Ctrl
APC Goat anti-mouse IgG (minimal x-reactivity)
PE Goat anti-mouse IgG (minimal x-reactivity)
Cell Staining Buffer

Clone
MOPC-21
Poly4053
Poly4053

Application
FC, ICFC, ICC, IF, IHC, IP, WB
FC
FC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.