

Product Data Sheet

Purified anti-NPM/B23-Phosphorylated (Thr199)

Catalog # / Size: 619201 / 50 µl (5 Western blots)

Clone: Poly6192

Isotype: Rabbit IgG


Immunogen: Modified peptide

Reactivity: Human, reacts against Thr199-phosphorylated NPM/B23

Preparation: The antibody was purified by antigen-affinity chromatography.

Formulation: This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 50% glycerol. Bovine serum albumin is present at 1 mg/ml final concentration (USA origin).

Storage: Upon receipt, store frozen at -20° C.



Hela cells were treated with 300 µM mimosine for 16 h, then placed in complete media (lane 1) or media containing 200 ng/ml nocodazole (lane 2) for an additional 18 h. Nuclear extract was western blotted with Poly6192. Proteins were visualized using a donkey anti-rabbit secondary conjugated to HRP and a chemiluminescence detection system.

Applications:


Applications: WB - Quality tested
IHC - Validated

Recommended Usage: Each lot of this antibody is quality control tested by Western blotting. Western blotting, suggested working dilution(s): Use 10 µl per 5 ml antibody dilution buffer for each mini-gel. For IHC, use a 1:50 dilution of antibody for staining. Antigen retrieval using 0.01 M sodium citrate buffer is recommended. It is recommended that the reagent be titrated for optimal performance for each application.

Description: NPM/B23 (also known as numatrin, nucleophosmin 1, nucleolar phosphoprotein B23) is a member of the nucleophosmin/nucleoplasmin family that shuttles between the nucleus and cytoplasm. This protein regulates the stability and transcriptional activity of p53 and acts as a molecular chaperone. NPM/B23 preferentially binds to denatured proteins and has been shown to stimulate DNA polymerase activity and control the duplication of centrosomes. The chaperone activity of NPM/B23 is regulated by protein kinase CK2 and promotes release of denatured substrates from NPM/B23. NPM/B23 is modified by phosphorylation on Thr199. This protein has been shown to bind to RNA and DNA and interact with nucleolar proteins including nucleolin, protein P120, HIV-1 Rev protein, and hepatitis delta antigens. The Poly6192 antibody recognizes human phosphorylated NPM/B23 (Thr199) and has been shown to be useful for Western blotting.

Antigen References: 1. Szebeni A, *et al.* 2003. *J. Biol. Chem.* 278:9107.
2. Colombo E, *et al.* 2002. *Nat. Cell Biol.* 4:529.
3. Chan PK, *et al.* 1986. *J. Biol. Chem.* 261:14335.

Related Products:	Product	Clone	Application
	Purified anti-Nucleolin-Phosphorylated (Thr76/Thr84)	10C7	WB, IHC
	Purified anti-NPM/B23-Phosphorylated (Thr234/Thr237)	Poly6191	WB, IHC
	HRP Donkey anti-rabbit IgG (minimal x-reactivity)	Poly4064	ELISA, IHC, WB



Formalin-fixed paraffin-embedded human thymus tissue was stained with Poly6192 and developed with an alkaline phosphatase chromogen substrate (red color). Tissue was counterstained with H&E (blue/pink). Magnification, 40X.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.