

# Product Data Sheet

## Purified anti-rat IFN- $\gamma$

**Catalog # / Size:** 507801 / 50  $\mu$ g  
507802 / 500  $\mu$ g

**Clone:** DB-1

**Isotype:** Mouse IgG1,  $\kappa$

**Immunogen:** Recombinant rat IFN- $\gamma$


**Reactivity:** Mouse, Rat

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



## Applications:

**Applications:** ELISA Capture - *Quality tested*  
ICFC, IHC, WB - *Reported in the literature*

**Recommended Usage:** Each lot of this antibody is quality control tested by ELISA analysis. For ELISA capture applications, the suggested use of this reagent is 1-4  $\mu$ g/ml. To obtain a linear standard curve, serial dilutions of IFN- $\gamma$  recombinant protein ranging from 1000 to 8 pg/ml are recommended for each ELISA plate. The purified DB-1 has been tested by blocking fluorochrome conjugated DB-1 for intracellular cytokine staining. In order to obtain complete blocking results, a saturated amount of purified antibody ( $\leq 5.0$   $\mu$ g/million cells) should be used for incubation with target cells, prior to staining with fluorochrome conjugated antibody. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** **ELISA Capture<sup>1</sup> or ELISPOT Capture<sup>2</sup>:** The purified DB-1 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated poly5109 antibody (Cat. No. 510901) as the detecting antibody and recombinant IFN- $\gamma$  (Cat. No. 565701) as the standard. The LEAF™ purified antibody is suggested for ELISPOT capture.

**Flow Cytometry<sup>5</sup>:** The fluorochrome-labeled DB-1 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN- $\gamma$ -producing cells within mixed cell populations. For intracellular cytokine staining protocol, please visit [www.biolegend.com](http://www.biolegend.com) and click on the support section.

**Neutralization<sup>3,4</sup>:** The LEAF™ purified antibody (Endotoxin  $<0.1$  EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for neutralization of rat IFN- $\gamma$  bioactivity *in vivo* and *in vitro* (Cat. No. 507808).

**Additional reported applications (for the relevant formats) include:** Western blotting<sup>1</sup>, and immunohistochemistry<sup>2</sup> of paraformaldehyde-fixed, saponin-treated frozen tissue sections.

**Application References:**

1. Van der Meide P, *et al.* 1989. *Lymphokine Res.* 8:439.
2. Nennesmo I, *et al.* 1989. *Brain Res.* 504:306.
3. Rayner D, *et al.* 1987. *Scand. J. Immunol.* 25:621.
4. Hartung H, *et al.* 1990. *Ann Neurol.* 27:247.
5. Bernard I, *et al.* 1998. *Eur. Cytokine Net.* 9:613.

**Description:** Interferon- $\gamma$  is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN- $\gamma$  also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- $\gamma$  can upregulate MHC class I and II antigen expression by antigen-presenting cells. The DB-1 antibody reacts with rat and mouse interferon-gamma (IFN- $\gamma$ ). The DB-1 antibody can neutralize the bioactivity of natural or recombinant IFN- $\gamma$ . The DB-1 antibody has been well characterized for ELISPOT, ELISA, intracellular staining, Western blotting, IHC, and neutralization (*in vitro* and *in vivo*).

**Antigen References:**

1. Fitzgerald K, *et al.* Eds. 2001. *The Cytokine FactsBook*. Academic Press San Diego.
2. De Maeyer E, *et al.* 1992. *Curr. Opin. Immunol.* 4:321.
3. Farrar M, *et al.* 1993. *Annu. Rev. Immunol.* 11:571.
4. Gray P, *et al.* 1987. *Lymphokines* 13:151.

### Related Products:

Product  
Purified Mouse IgG1,  $\kappa$  Isotype Ctrl  
TMB Substrate Reagent Set  
ELISA Assay Diluent (5X)  
ELISA Coating Buffer (5X)

Clone  
MOPC-21

Application  
FC, ICFC, ICC, IF, IHC, IP, WB  
ELISA  
ELISA  
ELISA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.