

## **Product Data Sheet**

102

Log Fluorescence Intensity

LPS-stimulated (6 hours, in the

presence of monensin) human peripheral blood monocytes

intracellularly stained with E8N1 Alexa Fluor® 488

103

104

cell Numbe

100

101

## Alexa Fluor® 488 anti-human IL-8

Catalog # / Size: 511411 / 25 tests

511412 / 100 tests

Clone: E8N1

**Isotype:** Mouse IgG1,  $\kappa$ 

Reactivity: Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 488 under optimal conditions. The solution is free of

unconjugated Alexa Fluor® 488.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.

## **Applications:**

Applications: ICFC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by intracellular

immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is 5 µl per 106 cells in 100 µl volume. It is recommended that the reagent be titrated for

optimal performance for each application.

\* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at

\*\* Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and

issued patents.

Application Notes: ELISA Detection: The biotinylated E8N1 antibody is useful as the detection antibody in a sandwich ELISA assay,

when used in conjunction with the purified H8A5 (Cat. No. 511502) antibody as the capture antibody. Flow Cytometry: The fluorochrome-labeled E8N1 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-8-producing cells within mixed cell populations. For intracellular cytokine staining

protocol, please visit www.biolegend.com and click on the support section.

Note: For testing human IL-8 in serum or plasma, BioLegend's ELISA Max™ Sets are specially developed and

recommended.

**Description:** IL-8, also known as neutrophil chemotactic factor, neutrophil activating protein, and monocyte-derived neutrophil chemotactic factor, is a member of the alpha (C-X-C) subfamily of chemokines called CXCL8. In response to proinflammatory stimuli, IL-8 is produced by monocytes, macrophages, T cells, neutrophils, and fibroblasts. IL-8 promotes neutrophil chemotaxis and degranulation. The 72 amino acid IL-8 is the predominant form secreted by monocytes and lymphocytes. The E8N1 antibody recognizes the human IL-8 protein and has been shown to be useful for intracellular immunofluorescence flow cytometric analysis and as ELISA detection antibody.

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.

2. Baggiolini M, et al. 1994. Adv. Immunol. 55:97.

3. Schröder J, et al. 1992. Immunology Ser. 57:387. 4. Zwahlen R, et al. 1993. Intl. Rev. Expt. Pathol. 34:27.

Related Products: Product

Cell Staining Buffer Fixation Buffer

Permeabilization Wash Buffer (10X)

Alexa Fluor® 488 Mouse IgG1, k Isotype Ctrl (ICFC)

Clone

MOPC-21

Application FC, ICC, ICFC ICC, ICFC ICC, ICFC, IHC

ICFC, IF



