

## **Product Data Sheet**

## PE anti-rat IL-4

Catalog # / Size: 511905 / 25 µg

511906 / 100 µg

Clone: OX-81

**Isotype:** Mouse IgG1,  $\kappa$ 

Immunogen: Recombinant rat IL-4

Reactivity: Rat

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution

is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not

freeze.

## **Applications:**

Applications: ICFC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric

analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10<sup>6</sup> cells in 100 µl

volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells and basophils. IL-4 elicits many different

biological responses, but has two dominant functions. The first is regulating differentiation of naïve CD4<sup>+</sup> T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10 and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated immune response controlled by Th1 cells. The second is regulating IgE and IgG1

production by B cells

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine Facts Book. Academic Press San Diego.

2. Boulay J, et al. 1992. Curr. Opin. Immunol. 4:294.

3. Dullens H, et al. 1991. In vivo 5:567.

4. Paul W. 1991. Blood 77:1859.

**Related Products: Product** PE Mouse IgG1, κ Isotype Ctrl (ICFC)

Cell Staining Buffer RBC Lysis Buffer (10X)

Permeabilization Wash Buffer (10X)

Clone MOPC-21 Application

ICFC FC, ICC, ICFC FC, ICFC





