

Product Data Sheet

Biotin anti-mouse IFN-γ

Catalog # / Size: 505704 / 500 µg

Clone: R4-6A2 **Isotype:** Rat IgG1, κ

Immunogen: Partially-purified, native mouse IFN-γ

Reactivity: Mouse


Preparation: The antibody was purified by affinity chromatography, and conjugated with

biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. Do not freeze.



Mouse IFN-γ Concentration (pg/mL)

Applications:

Applications: ELISA Detection, ELISPOT Detection

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For use as an ELISA detection antibody, a concentration range of 0.5-2.0 μg/ml is recommended. To obtain a linear standard curve, serial dilutions of IFN-γ

recombinant protein ranging from 2000 to 15 pg/ml are recommended for each ELISA plate.

For use as an ELISPOT detection antibody, a concentration range of 1-4 µg/ml is recommended. It is recommended

that the reagent be titrated for optimal performance for each application.

Application Notes: ELISA Capture^{1-4,6,10,11} or ELISPOT Capture⁵: The purified R4-6A2 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated XMG1.2 antibody (Cat. No. 505804) as the detecting antibody and recombinant mouse IFN-γ (Cat. No. 575309) as the standard. The LEAF™ purified antibody is suggested for ELISPOT capture.

ELISA or ELISPOT Detection: The biotinylated R4-6A2 antibody is useful as the detecting antibody in a sandwich

ELISA or ELISPOT Detection: The biotinylated R4-6A2 antibody is useful as the detecting antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the purified XMG1.2 antibody (Cat. No. 505802/505812) as the capture antibody and recombinant mouse IFN-γ (Cat. No. 575309) as the standard.

Neutralization^{1,2,9}: The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for neutralization of mouse IFN-γ bioactivity *in vivo* and *in vitro* (Cat. No. 505706).

Additional reported applications (for the relevant formats) include: immunohistochemical staining^{3,6,8} of

acetone-fixed frozen tissue sections and paraformaldehyde-fixed, saponin-treated frozen tissue sections.

Note: For testing mouse IFN-γ in serum, plasma or supernatant, BioLegend's ELISA Max™ Sets (Cat. No. 430801 to

430806) are specially developed and recommended.

1. Abrams, J., et al. 1992. Immunol. Rev. 127:5. Application References:

Stevenson, M., et al. 1990. Infect. Immun. 58:3225.

3. Sander, B., *et al.* 1993. *J. Immunol. Meth.* 166:201. 4. Yang, X., *et al.* 1993. *J. Immunoassay* 14:129.

5. Klinman, D., et al. 1994. Curr. Prot. Immunol.. John Wiley and Sons, New York. Unit 6.19. 6. Sander, B., et al. 1991. Immunol. Rev. 119:65.

7. Finkelman, F., et al. 2003. Curr. Prot. Immunol. John Wiley & Sons, New York. Unit 6.28.

8. Khanna, A., et al. 2000. J. Immunol. 164:1346.

9. Terrazas, L. I., *et al.* 2005. *Intl. J. Parasitology.* 35:1349. 10. Dzhagalov, I., *et al.* 2007. *J. Immunol.* 178:2113.

11. Xu, G., et al. 2007. J. Immunol. 179:5358.

Description: Interferon-y is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally

characterized based on anti-viral activities, IFN-γ also exerts anti-proliferative, immunoregulatory, and

proinflammatory activities. IFN-γ can upregulate MHC class I and II antigen expression by antigen-presenting cells. The R4-6A2 antibody reacts with mouse interferon-gamma (IFN-γ). The R4-6A2 antibody can neutralize the bioactivity

of natural or recombinant IFN-γ.

Antigen References: 1. Fitzgerald, K., et al. Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.

2. De Maeyer, E., *et al.* 1992. *Curr. Opin. Immunol.* 4:321. 3. Farrar, M., *et al.* 1993. *Annu. Rev. Immunol.* 11:571.

4. Gray, P., et al. 1987. Lymphokines 13:151.

Application **Related Products: Product** Clone

Purified anti-mouse IFN-γ XMG1.2 ELÍSA Capture, IHC, WB, CvTOF®

rm IFN-γ BA, ELISA Recombinant Mouse IFN-γ



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



HRP Avidin TMB Substrate Reagent Set ELISA Assay Diluent (5X) Mouse IFN-γ ELISA MAX™ Standard Mouse IFN-γ ELISA MAX™ Deluxe

Avidin

ELISA, ELISPOT, IHC, WB ELISA ELISA ELISA ELISA



