

Product Data Sheet

Purified anti-human IL-2

Catalog # / Size: 500301 / 50 µg

500302 / 500 µg

Clone: MQ1-17H12 **Isotype:** Rat IgG2a, κ

Immunogen: E. coli - expressed recombinant human IL-2

Reactivity: Human, Cross-Reactivity: Chimpanzee, Baboon, Cynomolgus, Rhesus,


Sooty Mangabey

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Applications:

Applications: ELISA Capture - Quality tested IHC, IP - Reported in the literature

CvTOF® - Validated

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration range of 2-6 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-2 recombinant protein

ranging from 2000 to 15 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated

for optimal performance for each application.

Application Notes: ELISA or ELISPOT Capture^{2,3}: The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich

ELISA or ELISPOT assay, when used in conjunction with the biotinylated Poly5176 antibody (Cat. No. 517601) as the detecting antibody. The LEAF™ purified antibody is suggested for ELISPOT capture. For ELISPOT capture

applications, a concentration range of 4-8 µg/ml is recommended.

Additional reported applications (for the relevant formats) include: immunoprecipitation², immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections^{1,4-6,8}, and immunocytochemistry.

Note: For testing human IL-2 in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 431801 to 431806) are

specially developed and recommended.

Application References: 1. Andersson J, et al. 1994. Immunology 83:16. (IHC)

2. Abrams J, et al. 1992. Immunol. Rev. 127:5. (ÎP) 3. Abrams JS. 1995. Curr. Prot. Immunol. Unit 6.20.

4. Fernandez V, et al. 1994. Eur. J. Immunol. 24:1808. (IHC)

5. Skansen-Saphir U, et al. 1994. Eur. J. Immunol. 24:916. (IHC)

Andersson U, et al. Detection and Quantification of Gene Expression. New York: Springer-Verlag. (IHC)

7. Prussin C, et al. 1995. J. Immunol. Methods. 188:117.

8. Raqib R, et al. 2002. Infect. Immun. 70:3199. (IHC)

9. Dzhagalov I, et al. 2007. J. Immunol. 178:2113. PubMed 10. Colleton BA, et al. 2009. J Virol. 83:6288. PubMed 11. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)

12. Rout N, et al. 2010. PLoS One 5:e9787. (FC)

Description: IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting

proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK

cells, LAK cells, monocytes, and oligodendrocytes.

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.

2. Taniguchi T, et al. 1993. Cell 73:5.

3. Nistico G. 1993. Prog. Neurobiol. 40:463.

4. Waldmann T, et al. 1993. Ann. NY Acad. Sci. 685:603.

Related Products: Product

Biotin anti-human IL-2 Recombinant Human IL-2

HRP Avidin

TMB Substrate Reagent Set ELISA Assay Diluent (5X)

Clone Poly5176 rh IL-2 Avidin

ELISA Detection BA, ELISA

ELÍSA, ELISPOT, IHC, WB **ELISA**

ELISA

Application



