

Technical Data Sheet

Recombinant Mouse IL-13

Product Information

Material Number:	554599
Size:	5 µg
Concentration:	100 µg/ml
Storage Buffer:	Frozen aqueous buffered solution containing BSA.

Description

Interleukin-13 (IL-13) is a cytokine produced by activated T lymphocytes which regulates the function of mouse macrophages and shares some biological functions with IL-4. Administration of mouse IL-13 to cultured mouse macrophages induces morphological changes (flattening, and development of processes), leads to increases in class II MHC antigen expression, and inhibits the synthesis of several proinflammatory cytokines (including IL-1β, IL-6, TNF, and IL-12). In contrast to human IL-13, mouse IL-13 does not appear to have activity for mouse B cells. Recombinant mouse IL-13 (mIL-13) has multiple glycosylated forms ranging in molecular weights from 11.3 to 15 kD, as detected by SDS-PAGE analysis.

Formulation and Purity:

Recombinant mouse IL-13 is supplied as a frozen aqueous buffered solution containing 1.0 mg/ml biotechnology grade, low endotoxin bovine serum albumin, with no preservatives. Recombinant mouse IL-13 is ≥95% pure, as determined by SDS-PAGE, and an absorbance assay based on the Beers- Lambert law. The endotoxin level is ≤0.1 ng per µg of mouse IL-13, as measured in a chromogenic LAL assay.

Preparation and Storage

Store product at -80°C prior to use or for long term storage of stock solutions.

Avoid multiple freeze-thaws of product.

Rapidly thaw and quick-spin product prior to use.

Upon initial thawing the product should be aliquoted into polypropylene microtubes and frozen at -80°C for future use.

Alternatively, prepare stock solutions diluted in sterile neutral buffer containing not less than 0.5 - 10 mg/ml carrier protein such as human or bovine serum albumin, aliquoted and stored at -80°C. For in vitro biological assay use, we recommend carrier-protein concentrations of 0.5 - 1 mg/ml. For use as an ELISA standard we recommend carrier-protein concentrations of 5 - 10 mg/ml. Failure to add carrier protein or store at indicated temperatures may result in a loss of activity. The product should not be diluted to less than 5 µg/ml for long term storage. Note: Carrier proteins should be pre-screened for possible effects in an appropriate experimental system. Carrier proteins may effect experimental results due to toxicity, high endotoxin levels or possible blocking activity.

Application Notes

Application

Bioassay	Routinely Tested
----------	------------------

Recommended Assay Procedure:

Biological Activity

Activity Assay: Measured by proliferation using TF-1 indicator cells

Specific Activity Range: 0.6 - 5 x 10⁷ Units*/mg

ED50: 0.2 - 1.5 ng/ml ; **Observed dose response** relationship: 50 fold

**Unit is defined as the amount of material required to stimulate a half-maximal response at cytokine saturation.*

Product Notices

1. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Cherwinski HM, Schumacher JH, Brown KD, Mosmann TR. Two types of mouse helper T cell clone. III. Further differences in lymphokine synthesis between Th1 and Th2 clones revealed by RNA hybridization, functionally monospecific bioassays, and monoclonal antibodies. *J Exp Med.* 1987; 166(5):1229-1244. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Doherty TM, Kastelein R, Menon S, Andrade S, Coffman RL.. Modulation of murine macrophage function by IL-13. *J Immunol.* 1993; 151(12):7151-7160. (Biology)

Kitamura T, Tange T, Terasawa T, et al. Establishment and characterization of a unique human cell line that proliferates dependently on GM-CSF, IL-3, or erythropoietin. *J Cell Physiol.* 1989; 140(2):323-334. (Methodology)

McKenzie A, Zurawski G. Measurement of IL-13. In: Coligan, Kruisbeek, Shevak, Strober, ed. *Current Protocols in Immunology*. New York: John Wiley & Sons; 1994:18-19. (Biology)

Zurawski G, de Vries JE.. Interleukin 13, an interleukin 4-like cytokine that acts on monocytes and B cells, but not on T cells. *Immunol Today.* 1994; 15(1):19-26. (Biology)