

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse CCL3/MIP-1 α in ELISAs and Western blots. In sandwich ELISAs, less than 0.03% cross-reactivity with recombinant human (rh) CCL3, recombinant mouse (rm) CCL9/10, and rmCCL4 and less than 0.4% cross-reactivity with rhCCL7.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse CCL3/MIP-1 α Ala24-Ala92 Accession # Q5QNW0
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Mouse CCL3/MIP-1 α Isoform LD78a (Catalog # 450-MA)
Immunohistochemistry	5-15 μ g/mL	Perfusion fixed frozen sections of mouse small intestine (Peyer's patch) and mouse thymus
Mouse CCL3/MIP-1α Sandwich Immunoassay		Reagent
ELISA Capture	0.2-0.8 μ g/mL	Mouse CCL3/MIP-1 α Antibody (Catalog # AF-450-NA)
ELISA Detection	0.1-0.4 μ g/mL	Mouse CCL3/MIP-1 α Biotinylated Antibody (Catalog # BAF450)
Standard		Recombinant Mouse CCL3/MIP-1 α (Catalog # 450-MA)
Neutralization	Measured by its ability to neutralize CCL3/MIP-1 α -induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR5. The Neutralization Dose (ND ₅₀) is typically 0.05-0.2 μ g/mL in the presence of 10 ng/mL Recombinant Mouse CCL3/MIP-1 α Isoform LD78a.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month from date of receipt, 2 to 8 °C, reconstituted. 6 months from date of receipt, -20 to -70 °C, reconstituted.

BACKGROUND

The macrophage inflammatory proteins 1 α and 1 β , two closely related but distinct proteins, were originally co-purified from medium conditioned by a LPS-stimulated murine macrophage cell line. Mature mouse MIP-1 α shares approximately 77% and 70% amino acid identity with human MIP-1 α and mouse MIP-1 β , respectively.

MIP-1 proteins are expressed primarily in T cells, B cells, and monocytes after antigen or mitogen stimulation. The MIP-1 proteins are members of the β (C-C) subfamily of chemokines.

Both MIP-1 α and MIP-1 β are monocyte chemoattractants *in vitro*. Additionally, the MIP-1 proteins have been reported to have chemoattractant and adhesive effects on lymphocytes, with MIP-1 α and MIP-1 β preferentially attracting CD8 $^{+}$ and CD4 $^{+}$ T cells, respectively. MIP-1 α has also been shown to attract B cells as well as eosinophils. MIP-1 proteins have been reported to have multiple effects on hematopoietic precursor cells and MIP-1 α has been identified as a stem cell inhibitory factor that can inhibit the proliferation of hematopoietic stem cells *in vitro* as well as *in vivo*. In the same assays, MIP-1 β was reported to be much less active. The functional receptor for MIP-1 α has been identified as CCR1 and CCR5.

References:

1. Menten, P. *et al.* (2002) Cytokine Growth Factor Rev. **13**:455.