



#### ORDERING INFORMATION

**Catalog Number:** AF1246

**Lot Number:** IRW01

**Size:** 100 µg

**Formulation:** 0.2 µm filtered solution in PBS  
with 5% trehalose

**Storage:** -20° C

**Reconstitution:** sterile PBS

**Specificity:** mouse 4-1BB Ligand

**Immunogen:** Sf21-derived rm4-1BB Ligand  
extracellular domain

**Ig Type:** mouse 4-1BB Ligand extracellular  
domain specific goat IgG

**Applications:** Direct ELISA  
Western blot

## ***Anti-mouse 4-1BB Ligand Antibody***

### ***Preparation***

Produced in goats immunized with purified, Sf21-derived, recombinant mouse 4-1BB Ligand (rm4-1BB Ligand) extracellular domain. Mouse 4-1BB Ligand specific IgG was purified by mouse 4-1BB Ligand affinity chromatography.

### ***Formulation***

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) with 5% trehalose.

### ***Endotoxin Level***

< 0.1 EU per 1 µg of the antibody as determined by the LAL method.

### ***Reconstitution***

Reconstitute with sterile PBS. If 1 mL of PBS is used, the antibody concentration will be 0.1 mg/mL.

### ***Storage***

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

### ***Specificity***

This antibody has been selected for its ability to recognize mouse 4-1BB Ligand in direct ELISAs and western blots.

### ***Applications***

**Direct ELISA** - This antibody can be used at 0.5 - 1.0 µg/mL with the appropriate secondary reagents to detect mouse 4-1BB Ligand. The detection limit for rm4-1BB Ligand is approximately 0.06 ng/well.

**Western blot** - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect mouse 4-1BB Ligand. The detection limit for rm4-1BB Ligand is approximately 1 ng/lane under non-reducing and reducing conditions. In this format, this antibody shows approximately 5% cross-reactivity with rmFas Ligand and less than 1% cross-reactivity with rmOX40 Ligand, rmTWEAK, rmTRAIL and rmTNF-α.

**Optimal dilutions should be determined by each laboratory for each application.**