

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human MICA in direct ELISAs and Western blots. In Western blots, approximately 40% cross-reactivity with recombinant human MICB is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MICA Ala23-Gln308 Accession # NP_001170990
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human MICA Fc Chimera (Catalog # 1300-MA)
Immunohistochemistry	5-15 µg/mL	See Below
Blockade of Receptor-ligand Interaction	In a functional ELISA, 0.4-1.2 µg/mL of this antibody will block 50% of the binding of 50 ng/mL of biotinylated Recombinant Human MICA Fc Chimera to immobilized Recombinant Human NKG2D Fc Chimera (Catalog # 1299-NK) coated at 2 µg/mL (100 µL/well). At 5 µg/mL, this antibody will block >90% of the binding.	

DATA

Immunohistochemistry



MICA in Human Breast Cancer Tissue. MICA was detected in immersion fixed paraffin-embedded sections of human breast cancer tissue using Goat Anti-Human MICA Antigen Affinity-purified Polyclonal Antibody (Catalog # AF1300) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Goat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS008) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of immersion fixed paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

MICA (MHC class I chain-related gene A) is a transmembrane glycoprotein that functions as a ligand for human NKG2D. A closely related protein, MICB, shares 85% amino acid identity with MICA. These proteins are distantly related to the MHC class I proteins. They possess three extracellular Ig-like domains, but they have no capacity to bind peptide or interact with β 2-microglobulin. The genes encoding these proteins are found within the Major Histocompatibility Complex on human chromosome 6. The MICA locus is highly polymorphic with more than 50 recognized human alleles. MICA is absent from most cells but is frequently expressed in epithelial tumors and can be induced by bacterial and viral infections. MICA is a ligand for human NKG2D, an activating receptor expressed on NK cells, NKT cells, $\gamma\delta$ T cells, and CD8⁺ $\alpha\beta$ T cells. Recognition of MICA by NKG2D results in the activation of cytolytic activity and/or cytokine production by these effector cells. MICA recognition is involved in tumor surveillance, viral infections, and autoimmune diseases.

References:

- Groh, V. *et al.* (2001) *Nature Immunol.* **2**:255.
- Stephens, H. (2001) *Trends Immunol.* **22**:378.
- Bauer, S. *et al.* (1999) *Science* **285**:727.
- Groh, V. *et al.* (2002) *Nature* **419**:734.
- Steinle, A. *et al.* (2001) *Immunogenetics* **53**:279.
- Pende, D. *et al.* (2002) *Cancer Res.* **62**:6178.
- NKG2D and its Ligands (2002) <http://www.RnDSystems.com>