

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PS-1 NTF in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinant human (rh) PS-1 CTF (aa 298-407) and rhPS-2 NTF (aa 1-80) is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Presenilin-1 Lys1-Lys80 Accession # P49768
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human Presenilin-1 N-Terminal Fragment
Immunohistochemistry	5-15 µg/mL	See Below

DATA

Immunohistochemistry



Presenilin-1 in Human Brain.

Presenilin-1 was detected in immersion fixed paraffin-embedded sections of human brain (hippocampus) using 15 µg/mL Goat Anti-Human Presenilin-1 N-Terminal Fragment Antigen Affinity-purified Polyclonal Antibody (Catalog # AF149) overnight at 4 °C. Tissue was stained with the Anti-Goat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS008) and counterstained with hematoxylin (blue). View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.