

Technical Data Sheet

FITC Mouse Anti-Mouse I-E[k]

Product Information

Material Number:	562015
Size:	50 µg
Concentration:	0.5 mg/ml
Clone:	14-4-4S
Immunogen:	C3H mouse skin graft and splenocytes
Isotype:	Mouse (C3H.SW) IgG2a, κ
Reactivity:	QC Testing: Mouse Reported: Rat
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The 14-4-4S antibody reacts with the I-E[k] MHC class II alloantigen. It cross-reacts with cells from mice of the H-2[d], H-2[p], and H-2[r] haplotypes. Cells from mice of the H-2[b], H-2[f], H-2[g7], H-2[q], and H-2[s] haplotypes do not express I-E antigen. It has been reported that mAb 14-4-4S cross-reacts with the rat MHC class II antigen RT1D.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.
The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.
The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
553456	FITC Mouse IgG2a, κ Isotype Control	0.25 mg	G155-178
561878	PE Rat Anti-Mouse CD45R/B220	25 µg	RA3-6B2
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
5. An isotype control should be used at the same concentration as the antibody of interest.

References

Blankenhorn EP, Symington FW, Cramer DV. Biochemical characterization of Ia antigens encoded by the RT1.B and RT1.D loci in the rat MHC. *Immunogenetics*. 1983; 17(5):475-484. (Clone-specific)
Hattori M, Buse JB, Jackson RA, et al. The NOD mouse: recessive diabetogenic gene in the major histocompatibility complex. *Science*. 1986; 231(4739):733-735. (Biology)
Klein J. Mutations in H-2E loci. In: Klein J. *Natural History of the Major Histocompatibility Complex*. New York: John Wiley & Sons; 1986:216-218. (Biology)
Neiss U, Reske K. Non-coordinate synthesis of MHC class II proteins and invariant chains by epidermal Langerhans cells derived from short-term in vitro culture. *Int Immunol*. 1994; 6(1):61-71. (Clone-specific)
Ozato K, Mayer N, Sachs DH. Hybridoma cell lines secreting monoclonal antibodies to mouse H-2 and Ia antigens. *J Immunol*. 1980; 124(2):533-540. (Immunogen)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD

