Technical Data Sheet

Purified Mouse Anti-Bcl-x

Product Information

Material Number:	551020			
Component:	51-6646GR			
Description:	Purified Mouse Anti- Bcl-x			
Size:	50 µg (1 ea)			
Concentration:	0.25 mg/ml			
Clone Name:	2H12			
Immunogen:	Human, mouse Bcl-xL / Bcl-xS (aa. 3-14)			
Isotype:	Mouse IgG2a, κ			
Target MW:	25-29 kDa			
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and $\leq 0.09\%$ sodium			
	azide.			
Component:	51-16526N			
Description:	Jurkat Cell Lysate			
Size:	50 µg (1 ea)			
Concentration:	1.0 mg/ml			
Storage Buffer:	SDS-PAGE buffer (62mM Tris pH 6.8, 2% SDS, 0.9% b-mercaptoethanol,			
	0.003% bromophenol blue, 5% glycerol)			

Description

Members of the Bcl-2 family play a major role in regulating the response of cells to a wide variety of apoptotic signals. Some Bcl-2 family members like Bcl-2, block apoptosis, whereas others promote apoptosis and inhibit Bcl-2 activity. Bcl-x is a Bcl-2 homologue that has two isoforms, resulting from alternative splicing. Bcl-xL (long) is a 241 amino acid protein (25-29 kDa) that is 47% homologous to Bcl-2 on the amino acid level. Bcl-xS (short) is a 178 amino acid protein (~ 19.5 kDa) lacking a 63 amino acid domain that is well conserved among members of the Bcl-2 protein family. Bcl-xL blocks cell death, whereas Bcl-xS inhibits Bcl-2 and promotes cell death.

The 2H12 antibody has been reported to recognize human, mouse and rat Bcl-xL (long) protein. An N-terminal peptide (amino acids 3-14) common to human and mouse Bcl-xL and Bcl-xS (short) proteins was used as the immunogen. Thus the antibody is also predicted to recognize the Bcl-xS protein as the sequence used for the immunogen is common to both Bcl-xL and Bcl-xS proteins. Reports from development, however, have indicated that Bcl-xS has not been observable.



Preparation and Storage

Store undiluted at -20°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

BD Biosciences

bd	bio	SCI	en	ces.	.com
	• •				~

bablobeleficeble	0111					
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean	
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995	
For country contact information, visit bdbiosciences.com/contact						
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD						



Application Notes

Application

Western blot	Routinely Tested
--------------	------------------

Recommended Assay Procedure:

Western blot: Please refer to http://www.bdbiosciences.com/support/resources/cell_biology/index.jsp

In addition to Jurkat cell lysates, MOLT-4 (Human T-lymphoblasts; ATCC CRL-1582), K-562 (Human bone marrow myelogenous leukemia; ATCC CCL-243), Hut-78 (Human T lymphoma; ATCC TIB-161), human peripheral blood mononuclear cells (PBMC), rat thymocytes and mouse thymocytes have also been reported to be useful as positive controls.

Suggested Companion Products

Catalog Number	Name	Size	Clone
611451	Jurkat Cell Lysate	500 μg	(none)
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.

- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 2.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 3. discarding to avoid accumulation of potentially explosive deposits in plumbing.

4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Hsu YT, Wolter KG, Youle RJ. Cytosol-to-membrane redistribution of Bax and Bcl-X(L) during apoptosis. Proc Natl Acad Sci U S A. 1997; 94(8):3668-3672. (Biology: Western blot)

Hsu YT, Youle RJ. Nonionic detergents induce dimerization among members of the Bcl-2 family. J Biol Chem. 1997; 272(21):13829-13834. (Immunogen: Western blot)

Krajewski S, Krajewska M, Shabaik A, et al. Immunohistochemical analysis of in vivo patterns of BcI-X expression. Cancer Res. 1994; 54(21):5501-5507. (Biology) Reed JC, Miyashita T, Krajewski S, et al. Bcl-2 family proteins and the regulation of programmed cell death in leukemia and lymphoma. Cancer Treat Res. 1996; 84:31-72. (Biology)

BD Biosciences

bdbiosciences.com United States

 Canada
 Europe
 Japan

 800.268.5430
 32.2.400.98.95
 0120.8555.90
Asia Pacific 877.232.8995 65.6861.0633 For country contact information, visit bdbiosciences.com/contact

For Country Contact Information, Visit **Dabiosciences.com/contact** Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

Latin America/Caribbean

55.11.5185.9995



