Technical Data Sheet FITC Mouse Anti-Human CD8

Product Information	
Material Number:	551347
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	G42-8
Isotype:	Mouse IgG2a, κ
Reactivity:	QC Testing: Human
Workshop:	NA
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide

Description

Reacts with the a subunit (32 kDa) of the two-chain complex. CD8 molecule binds to HLA class I molecules during interaction of CD8+T cells with antigen- presenting cells or with target cells. CD8 is expressed on T cytotoxic/ suppressor cell populations. G42-8 stains approximately 13-48% of peripheral blood lymphocytes and 80% of thymocytes, as well as a subset of NK cells. This antibody is useful in detecting CD8+ T cells when using DimerX I reagents (Cat. No. 551263).



Profile of peripheral blood lymphocytes analyzed by flow cytometry

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

A	Application					
	Flow cytometry	Routinely Tested				

Suggested Companion Products

Catalog Number	Name	Size	Clone
555573	FITC Mouse IgG2a, κ Isotype Control	100 tests	G155-178
551263	DimerX I: Recombinant Soluble Dimeric Human HLA-A2:Ig Fusion Protein	0.05 mg	(none)
BD Biosciences			

www.bdbiosciences.com								
United States Ca	nada	Europe	Japan	Asia Pacific	Latin America/Caribbean			
877.232.8995 88	8.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995			
For country-specific contact information, visit www.bdbiosciences.com/how_to_order/								
Conditions: The inform of any patents. BD Bio use of our products. Po product or as a compo	nation disclose sciences will no urchase does n onent of anoth	d herein is not to be ot be held responsil ot include or carry a er product. Any use	e construed as a reco ble for patent infring any right to resell or e of this product oth	ommendation to use gement or other vio r transfer this produce than the permitte	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express			

written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Schlossman S, Boumell L, et al, ed. Leucocyte Typing V. New York: Oxford University Press; 1995.(Biology)