

Technical Data Sheet


Purified Mouse Anti-Human Syk with Control

Product Information

Material Number:	551817
Size:	150 µg
Reactivity:	QC Testing: Human
Component:	51-16626N
Description:	EB1 Control Lysate
Size:	50 µg (1 ea)
Storage Buffer:	SDS-PAGE buffer (62mM Tris pH 6.8, 2% SDS, 0.9% b-mercaptoethanol, 0.003% bromophenol blue, 5% glycerol)
Component:	51-8116GR
Description:	Purified Mouse Anti-Human Syk
Size:	50 µg (3 ea)
Clone Name:	4D10
Immunogen:	Human Syk peptide aa. 314-339
Isotype:	Mouse IgG2a, κ
Storage Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09% sodium azide.

Description

Syk is a member of the family of non-receptor type protein-tyrosine kinases and plays a crucial role in lymphocyte signaling and development. Syk is expressed in all haematopoietic cells and contributes to the signal transduction process by binding to a tyrosine-based activation motif (ITAM) of immune receptors, including Igα, TCRζ, CD3ε, FcεRIβ, and FcεRIγ. Upon receptor activation, Syk binds to phosphorylated ITAMs via its two N-terminal SH2 domains thereby activating Syk and causing tyrosines in Syk to undergo auto-phosphorylation or phosphorylation. These phosphorylated sites can act as binding sites for other signaling molecules or help to regulate enzymatic activity. For example, in mast cells, Syk can activate downstream targets such as phospholipase Cγ1 and VAV. Studies in Syk-/- mast cells identified defects in both the ERK-MAP and JNK-MAP kinase pathways. Syk migrates as 72 kDa in SDS/PAGE. (Predicted MW from SWISS-PROT is 72 kDa; Accession number P43405).



Western blot analysis of Syk. Lysate from EB1 cells was probed with anti-Syk (clone 4D10) at concentrations of 0.125 (lane 1), 0.063 (lane 2), and 0.031 µg/ml (lane 3). Syk is identified as a band of ~72 kDa.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at -20°C.

Application Notes

Application

Western blot	Routinely Tested
--------------	------------------

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Recommended Assay Procedure:

Western Blot: EB1 control lysate [50 µg (1 µg/ml)] is provided as a positive control (store lysate at -20°C).

Additional application: The FITC-conjugated format of clone 4D10 is useful in immunofluorescent staining and flow cytometric analysis. Cat. No. 552476 is recommended for use in flow cytometry.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Chu DH, van Oers NS, Malissen M, Harris J, Elder M, Weiss A. Pre-T cell receptor signals are responsible for the down-regulation of Syk protein tyrosine kinase expression. *J Immunol.* 1999; 163(5):2610-2620.(Immunogen: Flow cytometry, Western blot)
Costello PS, Turner M, Walters AE, et al. Critical role for the tyrosine kinase Syk in signalling through the high affinity IgE receptor of mast cells. *Oncogene.* 1996; 13(12):2595-2605.(Biology)
Latour S, Veillette A. Proximal protein tyrosine kinases in immunoreceptor signaling. *Curr Opin Immunol.* 2001; 13:299-306.(Biology)
Turner M, Schweighoffer E, Colucci F, Di Santo JP, Tybulewicz VL. Tyrosine kinase SYK; essential functions for immunoreceptor signaling. *Immunol Today.* 2000; 21:148-154.(Biology)