

Technical Data Sheet

Purified Hamster Anti-Mouse TCR β Chain

Product Information

Material Number:	553167
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	H57-597
Immunogen:	TCR affinity-purified from mouse T-cell hybridoma DO-11.10
Isotype:	Armenian Hamster IgG2, λ 1
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The H57-597 antibody reacts with a common epitope of the β chain of the T-cell Receptor (TCR) complex on $\alpha\beta$ TCR-expressing thymocytes and peripheral T lymphocytes and NK1.1+ thymocytes and NK-T cells of all mouse strains tested. It does not react with $\gamma\delta$ TCR-bearing T cells. In the fetal and adult thymus, the TCR β chain may form homodimers or pair with the pre-TCR α chain on the surface of immature thymocytes before expression of the TCR. Plate-bound or soluble H57-597 antibody activates $\alpha\beta$ TCR-bearing T cells, and plate-bound mAb can induce apoptotic death.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4° C.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Reported
(Co)-stimulation	Reported
Cytotoxicity	Reported
Depletion	Reported
Western blot	Reported
Immunoprecipitation	Reported
Immunohistochemistry-formalin (antigen retrieval required)	Not Recommended

Recommended Assay Procedure:

It has been observed that pre-incubation of thymus cell suspensions at 37°C for 2 to 4 hours prior to staining enhances the ability of anti-CD3e and anti-TCR β chain mAbs to detect that T cell receptor on immature thymocytes.

Suggested Companion Products

Catalog Number	Name	Size	Clone
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
553030	FITC Rat Anti-Mouse CD8a	0.1 mg	53-6.7
554056	PE Mouse Anti-Armenian and Syrian Hamster IgG Cocktail	0.2 mg	(none)
553962	Purified Hamster IgG2, λ 1 Isotype Control	0.5 mg	Ha4/8

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



BD

BD Biosciences

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE™ (No Azide/Low Endotoxin) antibody format, if available, for in vitro and in vivo use.

References

- Atsuta N, Nishimura H, Nakamura N, Emoto M, Iwatsuki T, Yoshikai Y. Diversity of V gamma gene segments rearranged to the J gamma 4 gene in mice. *J Immunol.* 1995; 154(2):676-684.(Clone-specific: Depletion)
- Bendelac A, Killeen N, Littman DR, Schwartz RH. A subset of CD4+ thymocytes selected by MHC class I molecules. *Science.* 1994; 263(5154):1774-1778.(Biology)
- Bonneville M, Itoharu S, Krecko EG, et al. Transgenic mice demonstrate that epithelial homing of gamma/delta T cells is determined by cell lineages independent of T cell receptor specificity. *J Exp Med.* 1990; 171(4):1015-1026.(Clone-specific: Immunohistochemistry)
- Davenport C, Kumar V, Bennett M. Rapid rejection of H2k and H2k/b bone marrow cell grafts by CD8+ T cells and NK cells in irradiated mice. *J Immunol.* 1995; 155(8):3742-3749.(Clone-specific: Depletion)
- Duke RC, Cohen JJ, Boehme SA, et al. Morphological, biochemical, and flow cytometric assays of apoptosis. In: Coligan J, Kruisbeek AM, Margulies D, Shevach EM, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley and Sons; 1995:3.17.1-3.17.33.(Biology)
- Gascoigne NR. Transport and secretion of truncated T cell receptor beta-chain occurs in the absence of association with CD3. *J Biol Chem.* 1990; 265(16):9296-9301.(Clone-specific: Immunoprecipitation)
- Groettrup M, von Boehmer H. T cell receptor beta chain dimers on immature thymocytes from normal mice. *Eur J Immunol.* 1993; 23(6):1393-1396.(Biology)
- Kruisbeek AM, Shevach EM. Proliferative assays for T cell function. In: Coligan J, Kruisbeek AM, Margulies D, Shevach EM, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley and Sons; 1991:3.12.1-3.12.14.(Clone-specific: Stimulation)
- Kubo RT, Born W, Kappler JW, Marrack P, Pigeon M. Characterization of a monoclonal antibody which detects all murine alpha beta T cell receptors. *J Immunol.* 1989; 142(8):2736-2742.(Immunogen: Flow cytometry)
- Lefrancois L. Phenotypic complexity of intraepithelial lymphocytes of the small intestine. *J Immunol.* 1991; 147(6):1746-1751.(Biology)
- Ohno H, Ono S, Hirayama N, Shimada S, Saito T. Preferential usage of the Fc receptor gamma chain in the T cell antigen receptor complex by gamma/delta T cells localized in epithelia. *J Exp Med.* 1994; 179(1):365-369.(Clone-specific: Immunoprecipitation, Western blot)
- Saint-Ruf C, Panigada M, Azogui O, Debey P, von Boehmer H, Grassi F. Different initiation of pre-TCR and gammadeltaTCR signalling. *Nature.* 2000; 406(6795):524-527.(Clone-specific: Immunoprecipitation, Western blot)
- Saint-Ruf C, Ungewiss K, Groettrup M, Bruno L, Fehling HJ, von Boehmer H. Analysis and expression of a cloned pre-T cell receptor gene. *Science.* 1994; 266(5188):1208-1212.(Clone-specific: Stimulation)
- Skeen MJ, Ziegler HK. Induction of murine peritoneal gamma/delta T cells and their role in resistance to bacterial infection. *J Exp Med.* 1993; 178(3):971-984.(Clone-specific: Depletion)
- van der Heyde HC, Elloso MM, Chang WL, Kaplan M, Manning DD, Weidanz WP. Gamma delta T cells function in cell-mediated immunity to acute blood-stage Plasmodium chabaudi adami malaria. *J Immunol.* 1995; 154(8):3985-3990.(Clone-specific: Depletion)
- Vicari AP, Zlotnik A. Mouse NK1.1+ T cells: a new family of T cells. *Immunol Today.* 1996; 17(2):71-76.(Biology)
- Wagner DH Jr, Hagman J, Linsley PS, Hodsdon W, Freed JH, Newell MK. Rescue of thymocytes from glucocorticoid-induced cell death mediated by CD28/CTLA-4 costimulatory interactions with B7-1/B7-2. *J Exp Med.* 1996; 184(5):1631-1638.(Clone-specific: Cytotoxicity, Immunoprecipitation)