

Technical Data Sheet


Biotin Rat Anti-Mouse Vα2 TCR

Product Information

Material Number:	553287
Size:	0.25 mg
Concentration:	0.5 mg/ml
Clone:	B20.1
Immunogen:	Soluble αβ TCR from mouse cytotoxic T-cell clone KB5-C20
Isotype:	Rat (LOU) IgG2a, λ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The B20.1 antibody reacts with most members of the Vα2 T-cell Receptor (TCR) subfamily in mice having the *a*, *b*, and *c* haplotypes of the *Tcrb* gene complex. B20.1 antibody may cross-react with Vδ8 TCR, which shares >90% sequence homology with Vα2 TCR. Levels of B20.1+ T cells appear to be influenced by Vα haplotypes. Moreover, the frequencies of Vα2+ CD8+ and CD4+ T cells are influenced by H-2 haplotypes.



Two-color analysis of the expression of Vα2 TCR on peripheral lymphocytes. BALB/c lymph node cells were incubated simultaneously with biotinylated B20.1, PE-conjugated RM4-5 (anti-CD4, Cat. No. 553048/553049), and PE-conjugated 53-5.8 (anti-CD8b.2, Cat. No. 553041) monoclonal antibodies, followed by Avidin-FITC (Cat. No. 554057). Flow cytometry was performed on a FACScan™ (BDIS, San Jose, CA).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

For flow cytometry of cell suspensions from peripheral lymphoid tissues, it is recommended that multicolor staining be performed to distinguish T lymphocytes from non-T cells.

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
553048	PE Rat Anti-Mouse CD4	0.1 mg	RM4-5
553041	PE Rat Anti-Mouse CD8b.2	0.1 mg	53-5.8
554057	Avidin FITC	0.5 mg	(none)
553997	Biotin Rat IgG2a, λ Isotype Control	0.25 mg	B39-4

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Gregoire C, Rebai N, Schweisguth F, et al. Engineered secreted T-cell receptor alpha beta heterodimers. *Proc Natl Acad Sci U S A*. 1991; 88(18):8077-8081. (Clone-specific)

Pircher H, Rebai N, Groettrup M, et al. Preferential positive selection of V alpha 2+ CD8+ T cells in mouse strains expressing both H-2k and T cell receptor V alpha a haplotypes: determination with a V alpha 2-specific monoclonal antibody. *Eur J Immunol*. 1992; 22(2):399-404.(Immunogen)