

## Technical Data Sheet


## APC Rat Anti-Mouse CD117

## Product Information

<b>Material Number:</b>	553356
<b>Alternate Name:</b>	c-KIT; W; SCFR; Stem Cell Factor Receptor; Sl; Steel Factor Receptor; Ssm
<b>Size:</b>	0.1 mg
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	2B8
<b>Immunogen:</b>	Mouse Bone Marrow Mast Cells
<b>Isotype:</b>	Rat (WI) IgG2b, $\kappa$
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

## Description

The 2B8 monoclonal antibody specifically binds to CD117 (c-Kit), a transmembrane tyrosine-kinase receptor that is encoded by the *Kit* gene (formerly dominant white spotting, *W*). The c-Kit ligand (also known as steel factor, stem cell factor, and mast cell growth factor) encoded by the *Kitl* gene (formerly steel, *Sl*), is a co-mitogen for hematopoietic stem cells, myeloerythroid progenitors and a mast-cell differentiation factor. The *KitW* and *KitSl* mutant alleles have similar pleiotropic effects on the development of melanocytes, germ cells, and the hematopoietic system. In the adult bone marrow, CD117 is expressed on hematopoietic progenitor cells, including CD90 (Thy-1) low, TER-119-, CD45R/B220-, CD11b (Mac-1)-, Ly-6G (Gr-1)-, CD4-, CD8-, and Sca-1 (Ly-6A/E)+ multipotent hematopoietic stem cells, progenitors committed to myeloid and/or erythroid lineages, and precursors of B and T lymphocytes. This widespread expression of CD117 in hematopoietic precursors is consistent with the participation of c-Kit and its ligand in the regulation of several hematopoietic lineages. Intrathymic expression of c-Kit and c-Kit ligand suggest that CD117 is also involved in the regulation of some events during the development of T lymphocytes. CD117 is also expressed by mast cells and by dendritic cells found in the periaarteriolar lymphocyte sheaths (T-cell areas) of splenic white pulp. The mAb 2B8 reportedly does not block the action of c-Kit. This clone 2B8 had been reported that there was cross-reactivity with rat.



**Two-color analysis of the expression of CD117 on mouse bone marrow cells.** A single-cell suspension of BALB/c bone marrow was simultaneously stained with FITC Rat Anti-Mouse CD45R/B220 (Cat. Nos. 553087/553088, both panels) and either APC Rat IgG2b,  $\kappa$  Isotype Control (Left Panel; Cat. No. 553991) or APC Rat Anti-Mouse CD117 (Right Panel; Cat. No. 553356) monoclonal antibodies. Flow cytometry was performed on a BD FACSCalibur™ flow cytometry system.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## BD Biosciences

bdbiosciences.com

<b>United States</b>	<b>Canada</b>	<b>Europe</b>	<b>Japan</b>	<b>Asia Pacific</b>	<b>Latin America/Caribbean</b>
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

**Conditions:** The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD



## Suggested Companion Products

Catalog Number	Name	Size	Clone
553991	APC Rat IgG2b, $\kappa$ Isotype Control	0.1 mg	A95-1
553087	FITC Rat Anti-Mouse CD45R/B220	0.1 mg	RA3-6B2
553088	FITC Rat Anti-Mouse CD45R/B220	0.5 mg	RA3-6B2
554656	Stain Buffer (FBS)	500 ml	(none)

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
4. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.
5. An isotype control should be used at the same concentration as the antibody of interest.
6. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.

## References

Anderson DM, Lyman SD, Baird A, et al. Molecular cloning of mast cell growth factor, a hematopoietin that is active in both membrane bound and soluble forms. *Cell*. 1990; 63(1):235-243. (Biology)

Austen KF, Boyce JA. Mast cell lineage development and phenotypic regulation. *Leuk Res*. 2001; 25(7):511-518. (Biology)

Domen J, Weissman IL. Hematopoietic stem cells need two signals to prevent apoptosis; BCL-2 can provide one of these, Kit/c-Kit signaling the other. *J Exp Med*. 2000; 192(12):1707-1718. (Biology)

Fadini GP, Sartore S, Schiavon M, Albiero M, Baesso I, Cabrelle A, Agostini C, Avogaro A. Diabetes impairs progenitor cell mobilisation after hindlimb ischaemia-reperfusion injury in rats. *Diabetologia*. 2006; 49(12):3075-3084. (Clone-specific)

Godfrey DI, Zlotnik A. Control points in early T-cell development. *Immunol Today*. 1993; 14(11):547-553. (Biology)

Huang E, Nock K, Beier DR, et al. The hematopoietic growth factor KL is encoded by the Sl locus and is the ligand of the c-kit receptor, the gene product of the W locus. *Cell*. 1990; 63(1):225-233. (Biology)

Ikuta K, Weissman IL. Evidence that hematopoietic stem cells express mouse c-kit but do not depend on steel factor for their generation. *Proc Natl Acad Sci U S A*. 1992; 89(4):1502-1506. (Immunogen: Flow cytometry, Immunoprecipitation)

Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell*. 1997; 91(5):661-672. (Biology)

Lian Z, Toki J, Yu C, et al. Intrathymically injected hemopoietic stem cells can differentiate into all lineage cells in the thymus: differences between c-kit+ cells and c-kit- low cells. *Stem Cells*. 1997; 15(6):430-436. (Biology)

Mirmonsef P, Shelburne CP, Fitzhugh Yeatman C 2nd, Chong HJ, Ryan JJ. Inhibition of Kit expression by IL-4 and IL-10 in murine mast cells: role of STAT6 and phosphatidylinositol 3'-kinase. *J Immunol*. 1999; 163(5):2530-2539. (Biology)

Ogawa M, Matsuzaki Y, Nishikawa S, et al. Expression and function of c-kit in hemopoietic progenitor cells. *J Exp Med*. 1991; 174(1):63-71. (Biology)

Pulendran B, Lingappa J, Kennedy MK, et al. Developmental pathways of dendritic cells in vivo: distinct function, phenotype, and localization of dendritic cell subsets in FLT3 ligand-treated mice. *J Immunol*. 1997; 159(5):2222-2231. (Biology)

Rodewald HR, Kretzschmar K, Swat W, Takeda S. Intrathymically expressed c-kit ligand (stem cell factor) is a major factor driving expansion of very immature thymocytes in vivo. *Immunity*. 1995; 3(3):313-319. (Biology)

Valent P. The riddle of the mast cell: kit(CD117)-ligand as the missing link?. *Immunol Today*. 1994; 15(3):111-114. (Biology)

Zsebo KM, Wypych J, McNiece IK, et al. Identification, purification, and biological characterization of hematopoietic stem cell factor from buffalo rat liver-conditioned medium. *Cell*. 1990; 63(1):195-201. (Biology)

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

<b>United States</b>	<b>Canada</b>	<b>Europe</b>	<b>Japan</b>	<b>Asia Pacific</b>	<b>Latin America/Caribbean</b>
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

