

## Technical Data Sheet

# FITC Mouse IgG2a, κ Isotype Control

### Product Information

<b>Material Number:</b>	553456
<b>Size:</b>	0.25 mg
<b>Concentration:</b>	0.5 mg/ml
<b>Clone:</b>	G155-178
<b>Immunogen:</b>	TNP-keyhole limpet hemocyanin
<b>Isotype:</b>	Mouse (BALB/c) IgG2a, κ
<b>Storage Buffer:</b>	Aqueous buffered solution containing ≤0.09% sodium azide.

### Description

The G155-178 clone has an unknown specificity. Trinitrophenal (TNP), the immunogen, is a hapten not expressed on human, mouse, rat or non-human primate cells. In the absence of specific binding, this antibody may bind non-specifically to Fc receptors. The immunoglobulin from clone G155-178 was selected as an isotype control following screening for low background on a variety of mouse and human tissues.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

### Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

### Application Notes

#### Application

Flow cytometry	Routinely Tested
Isotype control	Routinely Tested
Intracellular staining (flow cytometry)	Not Recommended

### Recommended Assay Procedure:

An isotype control should be used at the same concentration as the antibody of interest (e.g., ≤1 µg/million cells for flow cytometry). Cat. No. 554647 is the recommended isotype control for intracellular staining.

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554647	FITC Mouse IgG2a, κ Isotype Control	0.1 mg	G155-178

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

### BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States 877.232.8995    Canada 888.259.0187    Europe 32.53.720.550    Japan 0120.8555.90    Asia Pacific 65.6861.0633    Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



# BD

## BD Biosciences