Technical Data Sheet

FITC Mouse Anti-Mouse H-2K[d]

Product Information

Material Number: 553565 Size: 0.5 mg 0.5 mg/mlConcentration: SF1-1.1 Clone:

BALB/c mouse cells Immunogen: Mouse (SJL) IgG2a, κ Isotype: QC Testing: Mouse Reactivity:

Storage Buffer: Aqueous buffered solution containing protein stabilizer and ≤0.09% sodium

Description

The SF1-1.1 antibody reacts with the α3 domain of the H-2K[d] MHC class I alloantigen. Reactivity with other haplotypes (e.g., b, j, k, p, q, s, v) has not been observed. It has been reported that plate-bound SF1-1.1 mAb moderately enhances the apoptotic response of thymocytes to plate-bound 145-2C11 mAb (anti-mouse CD3e, Cat. No. 557306/553058).

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze

Application Notes

Application

Flow cytometry Routinely Tested

Suggested Companion Products

Catalog Number	Name	Size	Clone
553456	FITC Mouse IgG2a, κ Isotype Control	0.25 mg	G155-178

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

Abastado JP, Casrouge A, Kourilsky P. Differential role of conserved and polymorphic residues of the binding groove of MHC class I molecules in the selection of peptides. J Immunol. 1993; 151(7):3569-3575. (Clone-specific)

Noun G, Reboul M, Abastado JP, Jaulin C, Kourilsky P, Pla M. Alloreactive monoclonal antibodies select Kd molecules with different peptide profiles. J Immunol. 1996; 157(6):2455-2461. (Clone-specific)

Zhao Y, Iwata M. Cross-linking of the TCR-CD3 complex with CD4, CD8 or LFA-1 induces an anti-apoptotic signal in thymocytes: the signal is canceled by FK506. Int Immunol. 1995; 7(9):1387-1396. (Biology)

BD Biosciences

bdbiosciences.com

United States Canada Asia Pacific Europe 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD

