

Technical Data Sheet

Purified Mouse Anti-Mouse H-2K[b]/H-2D[b]

Product Information

Material Number:	553575
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	28-8-6
Immunogen:	C3H.SW mouse splenocytes
Isotype:	Mouse (C3H) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The 28-8-6 antibody reacts with the H-2K[b] and H-2D[b] MHC class I alloantigens. The reaction with H-2K[b] has been reported to be stronger than that with H-2D[b]. Its recognition of H-2D[b] is dependent upon the presence of $\beta 2$ microglobulin. Cross-reactivity with an epitope on the N-terminal domains, $\alpha 1$ and $\alpha 2$, of the H-2D[d] MHC class I alloantigen has also been described. Reactivity with other haplotypes (e.g., *f*, *k*, *p*, *q*, *r*, *s*) has not been observed.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

Application Notes

Application

Flow cytometry	Routinely Tested
Cytotoxicity	Reported

Recommended Assay Procedure:

For immunohistochemical staining of cells expressing MHC class I antigen of the *b* haplotype, we recommend the use of biotinylated anti-mouse H-2K[b] mAb AF6-88.5 in our special formulation for immunohistochemistry, Cat. no. 550550.

Suggested Companion Products

Catalog Number	Name	Size	Clone
550550	Biotin Mouse Anti-Mouse H-2K[b]	1.0 ml	AF6-88.5
553454	Purified Mouse IgG2a κ Isotype Control	0.5 mg	G155-178
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Allen H, Fraser J, Flyer D, Calvin S, Flavell R. Beta 2-microglobulin is not required for cell surface expression of the murine class I histocompatibility antigen H-2Db or of a truncated H-2Db. *Proc Natl Acad Sci U S A*. 1986; 83(19):7447-7451.(Biology)
 Evans GA, Margulies DH, Shykind B, Seidman JG, Ozato K. Exon shuffling: mapping polymorphic determinants on hybrid mouse transplantation antigens. *Nature*. 1982; 300(5894):755-757.(Biology)
 Ozato K, Sachs DH. Monoclonal antibodies to mouse MHC antigens. III. Hybridoma antibodies reacting to antigens of the H-2b haplotype reveal genetic control of isotype expression. *J Immunol*. 1981; 126(1):317-321.(Immunogen: Cytotoxicity)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD

