

## Technical Data Sheet

## Purified Rat Anti-Mouse CD62E

## Product Information

<b>Material Number:</b>	553749
<b>Alternate Name:</b>	E-selectin, ELAM-1
<b>Size:</b>	0.5 mg
<b>Concentration:</b>	0.5 mg/ml
<b>Clone:</b>	10E9.6
<b>Immunogen:</b>	Mouse brain capillary endothelioma bEnd.3 (TNF $\alpha$ -stimulated) Cell Line
<b>Isotype:</b>	Rat (LEW) IgG2a $\kappa$
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing $\leq$ 0.09% sodium azide.

## Description

The 10E9.6 antibody reacts with the 97-110 kDa cell surface glycoprotein E-selectin (CD62E), also known as endothelial-leukocyte adhesion molecule-1 (ELAM-1), which is expressed on endotoxin- or cytokine-stimulated mouse endothelial cells. A suspension of TNF $\alpha$  stimulated mouse brain capillary endothelioma cells, from the cell line bEnd.3, was used as the immunogen. The epitope recognized by mAb 10E9.6 has been mapped to the first and/or second complement regulatory protein repeat domains of E-selectin. The 10E9.6 antibody has been reported to block binding of a monocyte cell line to E-selectin in vitro and to block neutrophil migration in BALB/c, but not C57BL/6 mice. It has no effect on leukocyte rolling in TNF $\alpha$ -treated mouse venules or on in vitro adhesion of myeloid cells to E-selectin. Studies have demonstrated that Cutaneous Lymphocyte Antigen (CLA), recognized by mAb HECA-452 (Cat. no. 555946), may be a ligand for CD62E.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

## Preparation and Storage

Store undiluted at 4° C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

## Application Notes

## Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development
ELISA	Reported
Immunoprecipitation	Reported
Blocking	Reported

## Suggested Companion Products

Catalog Number	Name	Size	Clone
553927	Purified Rat IgG2a $\kappa$ Isotype Control	0.5 mg	R35-95

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

Borges E, Pendl G, Eytner R, Steegmaier M, Zollner O, Vestweber D. The binding of T cell-expressed P-selectin glycoprotein ligand-1 to E- and P-selectin is differentially regulated. *J Biol Chem.* 1997; 272(45):28786-28792.(Biology)

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



Bosse R, Vestweber D. Only simultaneous blocking of the L- and P-selectin completely inhibits neutrophil migration into mouse peritoneum. *Eur J Immunol.* 1994; 24(12):3019-3024.(Immunogen: Blocking, ELISA, Immunoprecipitation)

Eppihimer MJ, Wolitzky B, Anderson DC, Labow MA, Granger DN. Heterogeneity of expression of E- and P-selectins in vivo. *Circ Res.* 1996; 79(3):560-569. (Biology: Blocking)

Ley K, Bullard DC, Arbones ML, et al. Sequential contribution of L- and P-selectin to leukocyte rolling in vivo. *J Exp Med.* 1995; 181(2):669-675.(Biology)

Pendl GG, Robert C, Steinert M, et al. Immature mouse dendritic cells enter inflamed tissue, a process that requires E- and P-selectin, but not P-selectin glycoprotein ligand 1. *Blood.* 2002; 99(3):946-956.(Biology)

Ramos CL, Kunkel EJ, Lawrence MB, et al. Differential effect of E-selectin antibodies on neutrophil rolling and recruitment to inflammatory sites. *Blood.* 1997; 89(8):3009-3018.(Immunogen: Blocking)

Weller A, Isenmann S, Vestweber D. Cloning of the mouse endothelial selectins. Expression of both E- and P-selectin is inducible by tumor necrosis factor alpha. *J Biol Chem.* 1992; 267(21):15176-15183.(Biology)