

Technical Data Sheet

Purified Mouse Anti-Human Retinoblastoma Protein


Product Information

Material Number:	554144
Alternate Name:	RB
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	XZ55
Immunogen:	Recombinant carboxy-terminal human RB protein aa. 387-928
Isotype:	Mouse IgG1
Reactivity:	QC Testing: Human Reported: Mouse, Chicken, Xenopus, Quail
Target MW:	105-116 kDa
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The retinoblastoma gene encodes a nuclear phosphoprotein (Rb or p110[Rb]) which is expressed in most normal cells of vertebrates and acts as a tumor suppressor gene product. An underphosphorylated form of RB is mainly found in resting or fully differentiated cells, whereas the hyperphosphorylated form is present in proliferating cells. Only the underphosphorylated form of RB binds specifically to viral oncogenes such as SV40 large T, adenoviral E1A and HPV-E7. This interaction may partially contribute to the transforming activity of these viral oncoproteins. Rb also interacts with several cyclins including A, D, and E as well as the transcriptional activator E2F. The importance of these interactions for the biological function of Rb is still being elucidated.

XZ55 recognizes an epitope located between amino acids 443-622 of human Rb. It cross-reacts with chicken, mouse and Xenopus Rb, and a putative quail Rb. A recombinant carboxy-terminal (amino acids 387-928) human Rb protein was used as immunogen.



Western blot analysis of Rb in MOLT-4 human leukemia cell lysate. The XZ55 antibody (Cat. No. 554144) recognizes phosphorylated and underphosphorylated forms of Rb (~110-116 kDa).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

Application Notes

Application

Western blot	Routinely Tested
Immunoprecipitation	Reported
Gel shift	Reported

Recommended Assay Procedure:

Application includes western blot analysis (1-2 µg/ml). Rb migrates as multiple closely-spaced bands between approximately 110-116 kDa when sized on denaturing polyacrylamide gels (i.e. by SDS-PAGE). The different bands represent different Rb phosphorylation states, the higher molecular weight bands are more highly phosphorylated than the lower molecular weight bands. The level of phosphorylation is cell cycle

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



dependent, and may also be cell type dependent (not all forms are seen in all cell types that express Rb). Gel conditions influence the actual number of bands observed. In cases where optimal band separation is desired, use a 7 to 10% non-gradient long (≥ 12 inches) gel. MOLT-4 human leukemia cells (ATCC CRL-1582) are suggested as a positive control. Additional applications not routinely tested at BD Biosciences Pharmingen include immunoprecipitation (1-2 $\mu\text{g/ml}$) and electrophoretic mobility gel shift assays (EMSA).

Suggested Companion Products

Catalog Number	Name	Size	Clone
554002	HRP Goat Anti-Mouse Ig	1.0 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Guilhot C, Benchaibi M, Flechon JE, Samarut J. The 12S adenoviral E1A protein immortalizes avian cells and interacts with the avian RB product. *Oncogene*. 1993; 8(3):619-624.(Clone-specific: Western blot)

Hollingsworth RE Jr, Chen PL, Lee WH. Integration of cell cycle control with transcriptional regulation by the retinoblastoma protein. *Curr Opin Cell Biol*. 1993; 5(2):194-200.(Biology)

Hu QJ, Bautista C, Edwards GM, Defeo-Jones D, Jones RE, Harlow E. Antibodies specific for the human retinoblastoma protein identify a family of related polypeptides. *Mol Cell Biol*. 1991; 11(11):5792-5799.(Clone-specific: Immunoprecipitation, Western blot)

Livingston DM. Functional analysis of the retinoblastoma gene product and of RB-SV40 T antigen complexes. *Cancer Surv*. 1992; 12:153-160.(Biology)

Riley DJ, Lee EY, Lee WH. The retinoblastoma protein: more than a tumor suppressor. *Annu Rev Cell Biol*. 1994; 10:1-29.(Biology)

Shirodkar S, Ewen M, DeCaprio JA, Morgan J, Livingston DM, Chittenden T. The transcription factor E2F interacts with the retinoblastoma product and a p107-cyclin A complex in a cell cycle-regulated manner. *Cell*. 1992; 68(1):157-166.(Clone-specific: Gel shift)