

Technical Data Sheet

Purified Hamster Anti-Rat CD49b

Product Information

Material Number:	554998
Alternate Name:	Integrin $\alpha 2$ chain
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	Ha1/29
Immunogen:	Cultured Rat Glomerular Epithelial
Isotype:	Armenian Hamster IgG2, λ
Reactivity:	QC Testing: Rat Tested in Development: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The Ha1/29 antibody reacts with integrin $\alpha 2$ chain (CD49b or VLA-2 α chain), the 160-kDa transmembrane glycoprotein that non-covalently associates with the integrin $\beta 1$ subunit (CD29) to form the integrin $\alpha 2\beta 1$ complex known as VLA-2. VLA-2, a receptor for collagen and laminin, is expressed on activated lymphocytes, epithelial cells, and platelets. It has been reported that Ha1/29 antibody can block VLA-2 binding to collagen and laminin. Ha1/29 antibody also reacts with mouse CD49b.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4° C.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development
Immunoprecipitation	Reported
Immunocytochemistry (cytospins)	Reported
Blocking	Reported

Suggested Companion Products

Catalog Number	Name	Size	Clone
553962	Purified Hamster IgG2, $\lambda 1$ Isotype Control	0.5 mg	Ha4/8
554011	FITC Mouse Anti-Armenian and Syrian Hamster IgG Cocktail	0.5 mg	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

BD Biosciences

www.bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



References

Hemler ME. VLA proteins in the integrin family: structures, functions, and their role on leukocytes. *Annu Rev Immunol.* 1990; 8:365-400.(Biology)
Mendrick DL, Kelly DM. Temporal expression of VLA-2 and modulation of its ligand specificity by rat glomerular epithelial cells in vitro. *Lab Invest.* 1993; 69(6):690-702.(Immunogen: Flow cytometry, Immunocytochemistry (cytospins), Immunoprecipitation)
Mendrick DL, Kelly DM, duMont SS, Sandstrom DJ. Glomerular epithelial and mesangial cells differentially modulate the binding specificities of VLA-1 and VLA-2. *Lab Invest.* 1995; 72(3):367-375.(Clone-specific)

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

