

Technical Data Sheet


PE Rat Anti-Mouse CD3 Molecular Complex

Product Information

Material Number:	555275
Size:	0.2 mg
Concentration:	0.2 mg/ml
Clone:	17A2
Immunogen:	γδ TCR-positive T-T hybridoma D1
Isotype:	Rat (SD) IgG2b, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The 17A2 monoclonal antibody specifically binds to the T-cell receptor-associated CD3 complex that is expressed on many thymocytes and mature T lymphocytes. Plate-bound 17A2 antibody has been reported to induce IL-2 production by cultured T cells in the absence of accessory cells. The binding of the 17A2 antibody to T cells can be blocked by the anti-CD3e mAb 145-2C11. This suggests that the 17A2 antibody recognizes an epitope of the CD3 epsilon chain. In vivo treatment with 17A2 antibody has been reported to partially deplete T lymphocytes and temporarily down-modulates CD3 expression on T cells.



CD3 expression in spleen and thymus. BALB/c splenocytes were simultaneously stained with PE-conjugated mAb 17A2 and FITC-conjugated anti-mouse CD19 mAb 1D3 (Cat. No. 557398/553786, left panel). C3H/HeN thymocytes were stained with PE-conjugated 17A2 mAb (right panel, filled histogram) or unstained (right panel, empty histogram). Flow cytometry was performed on a BD FACScan™ flow cytometry system.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
553989	PE Rat IgG2b, κ Isotype Control	0.1 mg	A95-1

Product Notices

1. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.

BD Biosciences

bdbiosciences.com
United States 877.232.8995 Canada 888.268.5430 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 0800.771.7157
For country-specific contact information, visit bdbiosciences.com/how_to_order/
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



4. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.

References

Miescher GC, Schreyer M, MacDonald HR. Production and characterization of a rat monoclonal antibody against the murine CD3 molecular complex. *Immunol Lett.* 1989; 23(2):113-118. (Biology: Induction)

Mysliwicz J, Thierfelder S. Antilymphocytic antibodies and marrow transplantation. XII. Suppression of graft-versus-host disease by T-cell-modulating and depleting antimouse CD3 antibody is most effective when preinjected in the marrow recipient. *Blood.* 1992; 80(10):2661-2667. (Biology: Depletion)