

Technical Data Sheet

Purified Rat Anti-Mouse CD74

Product Information

Material Number:	555317
Alternate Name:	Ii
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	In-1
Immunogen:	A.TL mouse lymphocytes and B10.A mouse-derived CH1 B-cell lymphoma
Isotype:	Rat (WF) IgG2b, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The In-1 antibody reacts with CD74, the MHC class II-associated Invariant chain (Ii), expressed in I-A+ cell populations. Although it is predominantly expressed intracellularly, it has also been detected on the cell surface of some B-cell lines and at low levels on a small number of B lymphocytes. In the mouse, alternatively spliced transcripts encode 31-kDa and 41-kDa isoforms, and the cell-surface CD74 has a chondroitin sulfate side chain which allows it to interact with CD44. CD74 is directly involved in the intracellular transport of MHC class II molecules and antigen presentation, and it affects the maturation of T and B lymphocytes.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
Immunohistochemistry-frozen	Tested During Development
Immunohistochemistry-zinc-fixed	Tested During Development
Immunofluorescence	Reported
Immunoprecipitation	Reported
Western blot	Reported
Immunohistochemistry-formalin (antigen retrieval required)	Not Recommended

Suggested Companion Products

Catalog Number	Name	Size	Clone
554715	BD Cytotfix/Cytoperm Plus Kit (with BD GolgiStop)	250 tests	(none)
553986	Purified Rat IgG2b, κ Isotype Control	0.5 mg	A95-1
554016	FITC Goat Anti-Rat Ig	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Bertolino P, Rabourdin-Combe C. The MHC class II-associated invariant chain: a molecule with multiple roles in MHC class II biosynthesis and antigen presentation to CD4+ T cells. *Crit Rev Immunol*. 1996; 16(4):359-379.(Biology)

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Bikoff EK, Huang LY, Episkopou V, van Meerwijk J, Germain RN, Robertson EJ. Defective major histocompatibility complex class II assembly, transport, peptide acquisition, and CD4+ T cell selection in mice lacking invariant chain expression. *J Exp Med.* 1993; 177(6):1699-1712.(Clone-specific: Immunoprecipitation)

Bodmer H, Viville S, Benoist C, Mathis D. Diversity of endogenous epitopes bound to MHC class II molecules limited by invariant chain. *Science.* 1994; 263(5151):1284-1286.(Biology)

Cresswell P. Assembly, transport, and function of MHC class II molecules. *Annu Rev Immunol.* 1994; 12:259-293.(Biology)

Elliott EA, Drake JR, Amigorena S, et al. The invariant chain is required for intracellular transport and function of major histocompatibility complex class II molecules. *J Exp Med.* 1994; 179(2):681-694.(Clone-specific: Immunofluorescence)

Farr A, DeRoos PC, Eastman S, Rudensky AY. Differential expression of CLIP:MHC class II and conventional endogenous peptide:MHC class II complexes by thymic epithelial cells and peripheral antigen-presenting cells. *Eur J Immunol.* 1996; 26(12):3185-3193.(Clone-specific: Immunohistochemistry, Western blot)

Koch N, Koch S, Hämmerling GJ. Ia invariant chain detected on lymphocyte surfaces by monoclonal antibody. *Nature.* 1982; 299(5884):644-645.(Immunogen: Immunoprecipitation)

Mehring JH, Harris MR, Kindle CS, McCourt DW, Cullen SE. Characterization of fragments of the murine Ia-associated invariant chain. *J Immunol.* 1991; 146(3):920-927.(Clone-specific: Western blot)

Nakagawa T, Roth W, Wong P, et al. Cathepsin L: critical role in Ii degradation and CD4 T cell selection in the thymus. *Science.* 1998; 280(5362):450-453.(Clone-specific: Immunohistochemistry, Western blot)

Naujokas MF, Morin M, Anderson MS, Peterson M, Miller J. The chondroitin sulfate form of invariant chain can enhance stimulation of T cell responses through interaction with CD44. *Cell.* 1993; 74(2):257-268.(Clone-specific: Immunoprecipitation)

Shachar I, Flavell RA. Requirement for invariant chain in B cell maturation and function. *Science.* 1996; 274(5284):106-108.(Biology)

Viville S, Neefjes J, Lotteau V, et al. Mice lacking the MHC class II-associated invariant chain. *Cell.* 1993; 72(4):635-648.(Clone-specific: Immunohistochemistry)