

Technical Data Sheet


Biotin Mouse Anti-Human CD30

Product Information

Material Number:	555828
Alternate Name:	Ki-1
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	BerH8
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Workshop:	IV A105
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

Reacts with a 120 kD type I transmembrane glycoprotein expressed on in-vitro stimulated T and B cells. CD30 is an activation marker which was initially identified on Reed-Sternberg cells and is also expressed on a few extrafollicular T and B cells located at the rim of germinal centers. It is a cytokine receptor which belongs to the nerve growth factor receptor/tumor necrosis factor receptor (NGFR/TNFR) superfamily. CD30 interaction with CD30 ligand (CD30L) can mediate signals for proliferation, apoptosis and cytotoxicity of lymphoid cells.



Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
555747	Biotin Mouse IgG1 κ Isotype Control	100 tests	MOPC-21
554061	PE Streptavidin	0.5 mg	(none)

BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmlingen/protocols](http://www.bdbiosciences.com/pharmlingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

Schlossman SF, Boumsell L, Gilks W, et al, ed. *Leukocyte Typing V: White Cell Differentiation Antigens*. New York: Oxford University Press; 1995.(Biology)

Knapp W, Dorken B, Rieber EP, et al, ed. *Leucocyte Typing IV*. New York: Oxford University Press; 1989.(Clone-specific)

Bowen MA, Olsen KJ, Cheng L, Avila D, Podack ER. Functional effects of CD30 on a large granular lymphoma cell line, YT. Inhibition of cytotoxicity, regulation of CD28 and IL-2R, and induction of homotypic aggregation. *J Immunol*. 1993; 151(11):5896-5906.(Clone-specific)

Guesdon JL, Ternynck T, Avrameas S. The use of avidin-biotin interaction in immunoenzymatic techniques. *J Histochem Cytochem*. 1979; 27(8):1131-1139. (Biology)