

Technical Data Sheet

Purified Mouse Anti-GFAP


Product Information

| | |
|------------------|--|
| Material Number: | 556328 |
| Size: | 0.5 mg |
| Concentration: | 0.5 mg/ml |
| Clone: | 1B4 |
| Immunogen: | Cow spinal cord homogenate |
| Isotype: | Mouse IgG2b |
| Reactivity: | QC Testing: Rat Reported: Human, Mouse, Cow, Sheep, Dog, Pig, Rabbit, Guinea Pig, Chicken |
| Target MW: | 50 kDa |
| Storage Buffer: | Aqueous buffered solution containing ≤0.09% sodium azide. |

Description

GFAP (Glial Fibrillary Acid Protein) is the major protein of glial filaments in differentiated astrocytes. BD Pharmingen offers a panel of monoclonal antibodies (4A11, 1B4, 2E1) that specifically recognize GFAP. They do not cross-react with other intermediate filaments such as vimentin, neurofilament proteins, desmin, keratin, neurotubules or microfilaments. Bovine spinal cord homogenate was used as immunogen. This antibody has broad species reactivity, recognizing GFAP in brain homogenates from human, mouse, rat, cow, sheep, dog, pig, rabbit, guinea pig and chicken. 1B4 is particularly useful for identifying GFAP in immunohistochemistry of frozen and formalin-fixed, paraffin-embedded brain tissue sections. Additional applications include western blot analysis and indirect immunofluorescence of tissue-cultured cells.

The antibody is routinely tested by immunohistochemical staining of formalin-fixed, paraffin-embedded brain tissue samples, which were subjected to pretreatment with citrate or tissue unmasking fluid (see figure). Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Formalin-fixed, paraffin-embedded section of human brain stained for GFAP (clone 1B4, Cat. No. 556328) using a DAB chromogen and hematoxylin counterstain. Red arrows indicate astrocytes (positive).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4°C.

Application Notes

Application

| | |
|--|---------------------------|
| Immunohistochemistry-formalin (antigen retrieval required) | Routinely Tested |
| Immunofluorescence | Tested During Development |
| Western blot | Reported |
| Immunohistochemistry-frozen | Reported |

Recommended Assay Procedure:

BD Biosciences

bdbiosciences.com

| | | | | | |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| United States | Canada | Europe | Japan | Asia Pacific | Latin America/Caribbean |
| 877.232.8995 | 888.259.0187 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995 |

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Recommended Assay Procedure:

BD Pharmingen offers additional GFAP specific antibodies: clone 2E1 (Cat. No. 556329), 4A11 (Cat. No. 556327), and clones 1B4, 4A11, 2E1 combined and available as a "cocktail" (Cat. No. 556330). Applications include indirect immunofluorescence of tissue-cultured cells, immunohistochemical staining of formalin-fixed paraffin-embedded brain tissue sections (25 µg/ml); and western blot analysis (1-2 µg/ml). Rat brain is suggested as a positive control.

Suggested Companion Products

| Catalog Number | Name | Size | Clone |
|----------------|-----------------------------------|-----------|--------|
| 556329 | Purified Mouse Anti-GFAP | 0.5 mg | 2E1 |
| 556327 | Purified Mouse Anti-GFAP | 0.5 mg | 4A11 |
| 556330 | Purified Mouse Anti-GFAP Cocktail | 0.5 mg | (none) |
| 551011 | Anti-Mouse Ig HRP Detection Kit | 200 tests | (none) |

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

McLendon RE, Bigner DD. Immunohistochemistry of the glial fibrillary acidic protein: basic and applied considerations. *Brain Pathol.* 1994; 4(3):221-228. (Clone-specific)
McLendon RE, Burger PC, Pegram CN, Eng LF, Bigner DD. The immunohistochemical application of three anti-GFAP monoclonal antibodies to formalin-fixed, paraffin-embedded, normal and neoplastic brain tissues. *J Neuropathol Exp Neurol.* 1986; 45(6):692-703.(Clone-specific: Immunohistochemistry)
Pegram CN, Eng LF, Wikstrand CJ, McComb RD, Lee YL, Bigner DD. Monoclonal antibodies reactive with epitopes restricted to glial fibrillary acidic proteins of several species. *Neurochem Pathol.* 1985; 3(2):119-138.(Clone-specific: Immunohistochemistry, Western blot)