

## Technical Data Sheet


## PE Hamster Anti-Mouse CD11c

## Product Information

|                         |                                                                                         |
|-------------------------|-----------------------------------------------------------------------------------------|
| <b>Material Number:</b> | <b>557401</b>                                                                           |
| <b>Alternate Name:</b>  | Integrin $\alpha$ chain                                                                 |
| <b>Size:</b>            | 0.1 mg                                                                                  |
| <b>Concentration:</b>   | 0.2 mg/ml                                                                               |
| <b>Clone:</b>           | HL3                                                                                     |
| <b>Immunogen:</b>       | C57BL/6 Mouse Intestinal Intraepithelial Lymphocytes                                    |
| <b>Isotype:</b>         | Armenian Hamster IgG1, $\lambda$ 2                                                      |
| <b>Reactivity:</b>      | QC Testing: Mouse                                                                       |
| <b>Storage Buffer:</b>  | Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide. |

## Description

The HL3 antibody reacts with the integrin  $\alpha$  chain of gp150, 95 (CD11c/CD18) which is expressed on dendritic cells and CD4<sup>+</sup> CD8<sup>+</sup> intestinal intraepithelial lymphocytes (IEL) and is upregulated on IEL and lymph-node T cells following *in vivo* activation. CD11c is also found on human NK cells. Although its expression on mouse NK cells is not published, we have detected CD11c on mouse splenic NK cells. Cells of the monocyte/macrophage lineage have been reported to express low levels of CD11c. CD11c plays a role in binding of iC3b.



**Expression of CD11c on mouse spleen NK cells.** C57BL/6 splenocytes were stained simultaneously with FITC-conjugated anti-mouse NK-1.1 mAb PK136 (Cat. No. 553164) and either isotype control (left panel) or PE-conjugated mAb HL3 (right panel). Flow cytometry was performed on a FACScan™ flow cytometry system.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

|                |                  |
|----------------|------------------|
| Flow cytometry | Routinely Tested |
|----------------|------------------|

## Suggested Companion Products

| Catalog Number | Name                                         | Size   | Clone     |
|----------------|----------------------------------------------|--------|-----------|
| 553954         | PE Hamster IgG1, $\lambda$ 1 Isotype Control | 0.1 mg | G235-2356 |

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

| United States | Canada       | Europe        | Japan        | Asia Pacific | Latin America/Caribbean |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| 877.232.8995  | 888.259.0187 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995         |

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at [http://www.bdbiosciences.com/pharmingen/hamster\\_chart\\_11x17.pdf](http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf).

## References

- Barclay NA, Brown MH, Birkeland ML, et al, ed. *The Leukocyte Antigen FactsBook*. San Diego, CA: Academic Press; 1997. (Biology)
- Gao JX, Liu X, Wen J, et al. Differentiation of monocytic cell clones into CD8 alpha+ dendritic cells (DC) suggests that monocytes can be direct precursors for both CD8 alpha+ and CD8 alpha- DC in the mouse. *J Immunol*. 2003; 170(12):5927-5935. (Biology)
- Huleatt JW, Lefrancois L. Antigen-driven induction of CD11c on intestinal intraepithelial lymphocytes and CD8+ T cells in vivo. *J Immunol*. 1995; 154(11):5684-5693. (Immunogen: Immunoprecipitation)
- Larson RS, Springer TA. Structure and function of leukocyte integrins. *Immunol Rev*. 1990; 114:181-217. (Biology)
- Maraskovsky E, Brasel K, Teepe M, et al. Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J Exp Med*. 1996; 184(5):1953-1962. (Biology)
- Pulendran B, Lingappa J, Kennedy MK, et al. Developmental pathways of dendritic cells in vivo: distinct function, phenotype, and localization of dendritic cell subsets in FLT3 ligand-treated mice. *J Immunol*. 1997; 159(5):2222-2231. (Clone-specific: Immunohistochemistry)