

## Technical Data Sheet


## PE Mouse Anti-Human CD208

## Product Information

Material Number:	558126
Alternate Name:	DC-LAMP
Size:	100 tests
Vol. per Test:	20 µl
Clone:	110-1112
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

Reacts with CD208, also known as DC-LAMP or lysosome associate membrane protein. DC-LAMP is an intracellular molecule, of approximately 70 - 90 kDa, induced during dendritic cell (DC) maturation. It is undetected in immature DCs, but it is sharply expressed upon activation with LPS, TNFα, or CD154 in DCs from skin, tonsil, cultures of CD34+ cells in presence of GM-CSF and TNFα, and on monocyte cultures in the presence of GM-CSF and IL-4. Reports describe the transient expression of CD208 in the MHC class II compartment before the translocation of the MHC II molecules to the cell surface. It has been suggested that the biological role of DCLAMP could be its participation in peptide loading onto MHC class II.



*Profile of CD208, DC-LAMP (110-112) reactivity on fixed and permeabilized monocyte-derived cultured dendritic cells with 1% Paraformaldehyde and 70% Ethanol Fixation.*

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

## Suggested Companion Products

Catalog Number	Name	Size	Clone
555749	PE Mouse IgG1, κ Isotype Control	100 tests	MOPC-21

## BD Biosciences

www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

*Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.*

*For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



## Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100- $\mu$ l experimental sample (a test).
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

## References

Banchereau J, Briere F, Caux C. Immunobiology of dendritic cells. *Annu Rev Immunol.* 2000; 18:767-811.(Biology)  
de Saint-Vis B, Vincent J, Vandenabeele S. de Saint-Vis B, Vincent J, Vandenabeele S. *Immunity.* 1998; 9(3):325-336.(Biology)