

Technical Data Sheet

Alexa Fluor® 647 Mouse anti-Actin

Product Information


Material Number:	558624
Size:	100 tests
Vol. per Test:	5 µl
Clone:	C4/actin
Immunogen:	Chicken gizzard muscle Actin
Isotype:	Mouse IgG1, κ
Reactivity:	Tested: Human Reported: Dictyostelium discoideum, Physarum polycephalum Confirmed by western blot using purified antibody (Cat. No. 612656 or 612657): Chicken, Dog, Mouse, Rat, Green Algae Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Storage Buffer:

Description

Changes in cellular morphology, adhesion, and motility occur through the reorganization of the actin cytoskeleton. This reorganization of actin filaments results from interactions between actin and actin-binding proteins. Actin is a 42-kDa protein that is known as G-actin in its monomeric form. Polymerization of G-actin monomers leads to the generation of flexible filaments, 5-9 nm in diameter, called F-actin. F-actin may be organized in linear bundles called stress fibers or in two-dimensional networks. The latter are highly concentrated beneath the plasma membrane and form the actin cortex. Regulation of actin cytoskeletal dynamics occurs through actin-binding proteins. These proteins bind to G- and/or F-actin and regulate various aspects of actin cytoskeletal dynamics, such as polymerization and depolymerization of actin, cross-linking of actin filaments into bundles, interaction of actin-based structures with membranes and other cytoskeletal elements, and locomotion of actin-based structures. Thus, the actin cytoskeleton is a complex matrix consisting of G- and F-actin along with the multitude of interactions between these actin forms and a variety of different types of actin-binding proteins.

The C4 monoclonal antibody reacts with all known isoforms of actin in vertebrate muscle and non-muscle cells.



Immunofluorescent staining of human cell lines. U-2 OS cells (ATCC HTB-96) were cultured, fixed, permeabilized with cold methanol, stained with Alexa Fluor® 647 Mouse anti-Actin (pseudo-colored green), and counter-stained with Hoechst 33342 (pseudo-colored blue) according to the Recommended Assay Procedure. The images were captured on a BD Pathway™ 855 Bioimager System with a 20x objective and merged using BD Attovision™ software. This antibody also stains A549 (ATCC CCL-185) and HeLa (ATCC CCL-2) cells. Triton X-100 is not recommended as a permeabilization agent with this antibody.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Application Notes

Application

Bioimaging	Routinely Tested
------------	------------------

Recommended Assay Procedure:

- Seed the cells in appropriate culture medium at ~10,000 cells per well in a BD Falcon™ 96-well Imaging Plate (Cat. No. 353219), and culture overnight.
- Remove the culture medium from the wells, and fix the cells by adding 100 µl of fresh 3.7% Formaldehyde in PBS or BD Cytofix™ fixation buffer (Cat. No. 554655) to each well and incubating for 10 minutes at room temperature (RT).
- Remove the fixative from the wells, and permeabilize the cells by adding 100 µl of -20°C 90% methanol or -20°C BD™ Phosflow Perm Buffer III (Cat. No. 558050) to each well and incubating for 5 minutes at RT.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



4. Remove the permeabilizer, and wash the wells twice with 100 μ l of 1 $\times$ PBS.
5. Remove the PBS, and block the cells by adding 100 μ l of blocking buffer (3% FBS in 1 $\times$ PBS) or BD Pharmingen™ Stain Buffer (FBS) (Cat. No. 554656) to each well and incubating for 30 minutes at RT.
6. Remove the blocking buffer, dilute the antibody conjugate 1:10 in blocking buffer or Stain Buffer (FBS), and stain the cells by adding 50 μ l of the diluted antibody conjugate to each well and incubating for 1 hour at RT.
7. Remove the diluted antibody conjugate, and wash the wells three times with 100 μ l of 1 $\times$ PBS.
8. Remove the PBS, and counter-stain the nuclei by adding 100 μ l of a 2 μ g/ml solution of Hoechst 33342 (eg, Sigma-Aldrich Cat. No. B2261) in 1 $\times$ PBS to each well at least 15 minutes before imaging.
9. View and analyze the cells on an appropriate imaging instrument.

Suggested Companion Products

Catalog Number	Name	Size	Clone
558050	Perm Buffer III	125 ml	(none)
554655	Fixation Buffer	100 ml	(none)
554656	Stain Buffer (FBS)	500 ml	(none)
353219	BD Falcon™ 96-well Imaging Plate	1 box	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test when following the Recommended Assay Procedure. A Test is typically ~10,000 cells cultured in a well of a 96-well imaging plate.
2. Alexa Fluor is a registered trademark of Molecular Probes, Inc., Eugene, OR.
3. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Hanstein C, Lange U, Schneider-Poetsch HA, Grolig F, Wagner G. Detection of actin and localization of phytochrome in the green alga *Mougeotia* by monoclonal antibodies. *Acta Histochem Suppl.* 1991; 41:223-230.(Clone-specific)

Lessard JL. Two monoclonal antibodies to actin: one muscle selective and one generally reactive. *Cell Motil Cytoskeleton.* 1988; 10(3):349-362.(Immunogen)

Pantaloni D, Le Clairche C, Cartier M-F. Mechanism of Actin-Based Motility. *Science.* 2001; 292:1502-1506.(Biology)