

Technical Data Sheet


PE Mouse Anti-Human Fetal Hemoglobin

Product Information

Material Number:	560041
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	2D12
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

Reacts with fetal hemoglobin (HbF), a form of hemoglobin present during fetal development. HbF resembles adult hemoglobin (HbA) in possessing two alpha chains but differs in possessing two gamma instead of two beta chains. In normal adults, synthesis of HbF does persist at very low levels ($<1\%$ of total Hb) and is restricted to a small population of erythrocytes called F cells. Hemoglobin F-expressing erythrocytes are normally seen during pregnancy. An increase in the expression of fetal hemoglobin in adult peripheral red blood cells is a common feature in the genetic disorders of hemoglobin, sickle-cell disease (SCD) and beta thalassemia.



Profile of erythrocytes obtained from cord blood analyzed by flow cytometry

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

We recommend to use 0.05% cold glutaraldehyde at room temperature for 10 minutes to fix the cells, then use 0.1% Triton X-100 at room temperature for 10 minutes to permeabilize the cells.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554680	PE Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Campbell TA, Ware RE, Mason M. Detection of hemoglobin variants in erythrocytes by flow cytometry. *Cytometry*. 1999; 35(3):242-248. (Methodology)

Horiuchi K, Osterhout ML, Kamma H, Bekoe NA, Hirokawa KJ. Estimation of fetal hemoglobin levels in individual red cells via fluorescence image cytometry. *Cytometry*. 1995; 20(3):261-267. (Methodology)

Thorpe SJ, Thein SL, Sampietro M, Craig JE, Mahon B, Huehns ER. Immunochemical estimation of haemoglobin types in red blood cells by FACS analysis. *Br J Haematol*. 1994; 87(1):125-132. (Methodology)