

## Technical Data Sheet

## Alexa Fluor® 647 Mouse anti-Src (pY418)

## Product Information


Material Number:	560096
Alternate Name:	c-Src, p60-Src, SRC, SRC1
Size:	50 tests
Vol. per Test:	20 µl
Clone:	K98-37
Immunogen:	Phosphorylated Human Src Peptide
Isotype:	Mouse (BALB/c) IgG1, κ
Reactivity:	Confirmed: Human Predicted: Chicken, Mouse, Rat
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

## Description

The non-receptor protein tyrosine kinases of the Src family have similar domain structures with considerable sequence homologies. They are associated with cell membranes and are activated by a wide variety of cell-surface receptors. Through their participation in cell-signaling cascades that mediate an array of cellular responses, they control essential cellular activities.

Src, the prototype of the Src family, is encoded by the *SRC* protooncogene, which is a homolog of the *v-Src* gene of the Rous sarcoma virus. Src's oncogenic potential may be a consequence of its role as a regulator of cell growth, morphology, motility, and adhesion. The protein contains domains that mediate protein-protein interactions and regulation of its tyrosine kinase activity. Mechanisms that regulate Src activity include phosphorylations of serines and tyrosines in most of the domains and interactions with receptor proteins, integrins, and other binding proteins. In particular, autophosphorylation at tyrosine 418 (Y418) activates tyrosine kinase activity. Phosphorylations at other sites are mediated by kinases such as cdc2, CSK, and CHK in signaling cascades and may up- or down-regulate Src activity.

The K98-37 monoclonal antibody recognizes the phosphorylated Y418 in the protein kinase domain of activated Src. The orthologous phosphorylation sites in chicken, mouse, and rat Src are Y415, Y423, and Y418, respectively. Because of sequence similarities in this region, it may also recognize the homologous sites of some other Src family members: Lyn (pY396), Fyn (pY420), Hck (pY410), Lck (pY394), and Yes (pY425).



**Analysis of Src (pY418) in human T leukemia cells and peripheral blood lymphocytes.** Jurkat cells (ATCC TIB152, left panel) and human peripheral blood mononuclear cells (PBMC, middle panel) were either treated with 1 µM Staurosporine (EMD Biosciences, Cat. No. 569397) for 2 hours at 37°C (shaded histogram) or untreated (open histogram). The cells were fixed (BD Cytotfix™ buffer, Cat. No. 554655) for 10 minutes at 37°C, permeabilized with BD™ Phosflow Perm Buffer III (Cat. No. 558050) on ice for 30 minutes, and then stained with Alexa Fluor® 647 Mouse anti-Src (pY418). Lymphocytes were selected by their scatter profile during analysis of the PBMC. The data demonstrates that the level of phosphorylation of Src (pY418) decreases when protein kinase activity is inhibited by the treatment. Flow cytometry was performed on a BD™ FACSCalibur flow cytometry system.

The specificity of mAb K98-37 was confirmed by western blot analysis (right panel) using unconjugated Rabbit anti-Src (Cell Signaling Technology, Cat. No. 2123, left blot) and unconjugated Mouse anti-Src (pY418) (right blot) monoclonal antibodies on lysates of Jurkat cells that were untreated (lanes 1) or treated with Staurosporine for 1 hour (lanes 2) or 2 hours (lanes 3). Src (pY418) is identified as a band of 60 kDa that is reduced by Staurosporine treatment.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	A-431	none	Cytofix	Perm III	Positive expression
	Human	A-431	Phospho peptide	Cytofix	Perm III	Blocking
	Human	A-431	Non-phospho peptide	Cytofix	Perm III	No blocking
	Human	Jurkat	Staurosporine	Cytofix	Perm III	Down-regulation
	Human	PBMC	Staurosporine	Cytofix	Perm I, II, or III	Down-regulation
WB	Human	A-431	Phospho peptide			Blocking of 60-kDa band
	Human	A-431	Non-phospho peptide			No blocking of 60-kDa band
	Human	Jurkat	Staurosporine			60-kDa band down-regulated
	Human	PBMC	Staurosporine			60-kDa band down-regulated

## Application Notes

### Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

### Recommended Assay Procedure:

Either BD Cytofix™ fixation buffer or BD™ Phosflow Fix Buffer I may be used for cell fixation. Any of the three BD™ Phosflow permeabilization buffers may be used.

### Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 ml	(none)
557870	Fix Buffer I	250 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558052	Perm Buffer II	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)

### Product Notices

- Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100- $\mu$ l experimental sample (a test).
- Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
- For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
- The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.

### References

Alvarez RH, Kantarjian HM, Cortes JE. The role of Src in solid and hematologic malignancies. Development of new-generation Src inhibitors. *Cancer*. 2006; 107(8):1918-1929. (Biology)

Arias-Salgado EG, Lizano S, Sarkar S, Brugge JS, Ginsberg MH, Shattil SJ. Src kinase activation by direct interaction with the integrin  $\beta$  cytoplasmic domain. *Proc Natl Acad Sci U S A*. 2003; 100(23):13298-13302. (Biology)

Frame MC. Newest findings on the oldest oncogene; how activated src does it. *J Cell Sci*. 2004; 117(7):989-998. (Biology)