Technical Data Sheet Alexa Fluor® 555 Mouse anti-Human TRA-1-81 Antigen

Product Information	
Material Number:	560123
Size:	100 tests
Vol. per Test:	5 µl
Clone:	TRA-1-81
Immunogen:	Human Embryonal Carcinoma Cell Line
Isotype:	Mouse (BALB/c) IgM, ĸ
Reactivity:	Confirmed: Human Reported: Rhesus Monkey
Storage Buffer:	Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09%

Description

The TRA-1-81 monoclonal antibody reacts with a pluripotent-stem-cell-specific epitope on a high-molecular-weight transmembrane glycoprotein. The TRA-1-81 antigen is an epitope on the same keratan sulfate core molecule, podocalyxin, as 4 other distinct antigens on tumor-derived cell lines, TRA-1-60, GCTM2, K4, and K21. The expression of TRA-1-81 antigen is stage-specific and can be used to characterize embryonic cells and monitor their differentiation. The antigen is found on teratocarcinoma (embryonal carcinoma or EC), embryonic inner cell mass (but not morula or trophoblast), and embryonic stem (ES) cells. As human EC and ES cells undergo differentiation, expression of TRA-1-81 antigen is lost.

sodium azide.

This antibody is routinely tested by immunofluorescent imaging. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.





Immunofluorescent staining of human ES cell line. The H9 cell line (WiCell, Madison, WI) was cultured, fixed, and stained with Alexa Fluor® 555 Mouse anti-Human TRA-1-81 Antigen monoclonal antibody (pseudo-colored green) according to the Recommended Assay Procedure. The left image shows the plasma membrane staining by the TRA-1-81 mAb, and the right image shows TRA-1-81 with counter-staining of the nuclei by Hoechst 33342 (pseudo-colored blue). The images were captured on a BD Pathway™ 435 Cell Analyzer using a 10X objective and merged using BD Attovision™ software.

Preparation and Storage

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated to Alexa Fluor® 555 under optimum conditions, and unreacted Alexa Fluor® 555 was removed.

Application Notes

Application							
Bioimaging		Routinely Tested					
BD Bioscie	ences						
www.bdbiosci	iences.com						
United States 877.232.8995 For country-sp	Canada 888.259.0187 ecific contact in	Europe 32.53.720.550 formation, visit v	Japan 0120.8555.90 www.bdbioscien	Asia Pacific 65.6861.0633 ces.com/how_to	Latin America 55.11.5185.99 _order/	a/Caribbean 195	
Conditions: The in of any patents. Bu use of our product product or as a co	nformation disclose D Biosciences will n cts. Purchase does r omponent of anoth	d herein is not to b ot be held responsi not include or carry er product. Any us	e construed as a rec ble for patent infrin any right to resell or e of this product oth	ommendation to use gement or other vio r transfer this produ er than the permitt	e the above produ lations that may o ct either as a stand ed use without the	ct in violation ccur with the -lalone e express	

written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

Bioimaging Certified Reagent

Recommended Assav Procedure:

- Seed the cells in appropriate culture medium at an appropriate cell density in a BD Falcon[™] 96-well Imaging Plate (Cat. No. 353219), and 1. culture overnight to 48 hours.
- Remove the culture medium from the wells, wash the wells twice with 100 µl of 1× PBS, and fix the cells by adding 100 µl of fresh 3.7% 2. Formaldehyde in PBS or BD Cytofix™ fixation buffer (Cat. No. 554655) to each well and incubating for 10 minutes at room temperature (RT)
- 3 Remove the fixative from the wells, and wash the wells twice with 100 μ l of 1× PBS.
- 4. Dilute the antibody in $1 \times PBS$, and stain the cells by adding 50 µl of the diluted antibody conjugate to each well and incubating for 1 hour at RT.
- 5. Remove the diluted antibody, and wash the wells twice with 100 μ l of 1× PBS.
- Remove the PBS, and counter-stain the nuclei by adding 100 µl of a 2 µg/ml solution of Hoechst 33342 (eg, Sigma-Aldrich Cat. No. B2261) 6. in 1× PBS to each well at least 15 minutes before imaging.
- View and analyze the cells on an appropriate imaging instrument. 7.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 ml	(none)
353219	BD Falcon [™] 96-well Imaging Plate	1 box	(none)

Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test when following the Recommended Assay Procedure. A Test is typically ~10,000 cells cultured in a well of a 96-well imaging plate.
- Alexa Fluor is a registered trademark of Molecular Probes, Inc., Eugene, OR. 2.
- 3. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
- 4 Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- Source of all serum proteins is from USDA inspected abattoirs located in the United States. 5.

References

Andrews PW, Banting G, Damanov I, Arnaud D, Avner P. Three monoclonal antibodies defining distinct differentiation antigens associated with different high molecular weight polypeptides on the surface of human embryonal carcinoma cells. Hybridoma. 1984; 3(4):347-361.(Immunogen: Immunofluorescence, Immunoprecipitation, Radioimmunoassav)

Badcock G, Pigott C, Goepel J, Andrews PW. The human embryonal carcinoma marker antigen TRA-1-60 is a sialylated keratan sulfate proteoglycan. Cancer Res. 1999: 59:4715-4719.(Clone-specific: Immunoprecipitation, Western blot)

Draper JS, Pigott C, Thomson JA, Andrews PW. Surface antigens of human embryonic stem cells: changes upon differentiation in culture. J Anat. 2002; 200:249-258.(Clone-specific: Flow cytometry)

Henderson JK, Draper JS, Baillie HS, et al. Preimplantation human embryos and embryonic stem cells show comparable expression of stage-specific embryonic antigens. Stem Cells. 2002; 20:329-337.(Clone-specific: Flow cytometry, Immunofluorescence)

Schopperle WM, DeWolf WC. The TRA-1-60 and TRA-1-81 human pluripotent stem cell markers are expressed on podocalyxin in embryonal carcinoma. Stem Cells. 2007; 25:723-730.(Clone-specific: Immunofluorescence, Western blot)

Thomson JA, Itskovitz-Eldor J, Shapiro SS, et al. Embryonic stem cell lines derived from human blastocysts. Science. 1998; 282:1145-1147. (Clone-specific: Immunocytochemistry (cytospins))

Thomson JA, Kalishman J, Golos TG, et al. Isolation of a primate embryonic stem cell line. Proc Natl Acad Sci U S A, 1995; 92:7844-7848.(Clone-specific: Immunocytochemistry (cytospins))

BD Biosciences

www.bdbiosciences.com								
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean			
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995			
For country-specific contact information, visit www.bdbiosciences.com/how_to_order/								
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation								
of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the								
use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone								
product or as a component of another product. Any use of this product other than the permitted use without the express								



written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD