

Technical Data Sheet

Alexa Fluor® 647 Mouse anti-eIF4E (pS209)



Product Information

Material Number:	560240
Alternate Name:	eIF-4E, CBP, EIF4E1, EIF4EL1, EIF4F, mRNA cap-binding protein
Size:	50 tests
Vol. per Test:	20 µl
Clone:	J77-925
Immunogen:	Phosphorylated Human eIF4E Peptide
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human Predicted: Mouse, Rat
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The eukaryotic translation *Initiation Factor 4E* (eIF4E) is a 25-kDa phosphoprotein that specifically binds to the 7-methylguanosine-containing cap of mRNA. eIF4E is the rate-limiting component for the initiation of cap-dependent translation by the eIF4F translation initiation complex, which is composed of eIF4E, eIF4G, and eIF4A. This complex promotes the unwinding of secondary structure at the 5' untranslated region of mRNA that is necessary to expose and locate the AUG initiation codon. Other functions of eIF4E have been identified, such as promoting the export of mRNAs that are involved in cell cycle progression from the nucleus and differentially regulating the translation of certain mRNAs in the cytoplasm. Three mechanisms for eIF4E regulation have been identified: Mnk1-mediated phosphorylation on serine 209 (S209) is required for eIF4E binding to the cap structure; over-expression of phosphorylated eIF4E can lead to increased cell proliferation, suppression of apoptosis, and a transformed phenotype; and interactions with nonphosphorylated eIF4E-binding proteins inhibit the formation of the eIF4F complex.

The J77-925 monoclonal antibody recognizes the phosphorylated S209 (pS209) of eIF4E.



Analysis of eIF4E (pS209) in monocytes. Human peripheral blood mononuclear cells (PBMC) were either stimulated with 40 nM Phorbol 12-myristate 13-acetate (PMA, Sigma-Aldrich Cat. No. P8139) for 15 minutes (shaded histogram) or unstimulated (open histogram). The cells were fixed (BD Cytofix™ buffer, Cat. No. 554655) for 10 minutes, then permeabilized (BD™ Phosflow Perm Buffer II, Cat. No. 558052) on ice for at least 30 minutes, and then stained with Alexa Fluor® 647 Mouse anti-eIF4E (pS209). Monocytes were selected by scatter profile. Flow cytometry was performed on a BD FACSAArray™ bioanalyzer system.

The specificity of mAb J77-925 was confirmed by western blot analysis using unconjugated antibody on lysates from control (lane 1) and PMA-treated (lane 2) PBMC. eIF4E (pS209) is identified as a band of 25 kDa, with increased intensity in the treated cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to Alexa Fluor® 647 under optimum conditions, and unreacted Alexa Fluor® 647 was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



The purified or conjugated mAb was characterized by flow cytometry (Flow) and western blot (WB) using these model systems:

Method	Species	Cells	Treatment	Fixation	Perm buffer	Result
Flow	Human	HeLa S3	Anisomycin	Cytofix	Perm III	Unsatisfactory
	Human	HeLa S3	λ or alkaline phosphatase	Cytofix	Perm III	Unsatisfactory
	Human	Jurkat	PMA	Cytofix	Perm III	Induction observed
	Human	Whole Blood	PMA	Lyse/Fix	Perm III	Unsatisfactory
	Human	PBMC	Anisomycin	Cytofix	Perm III	Unsatisfactory
	Human	PBMC	PMA	Lyse/Fix	Perm III	Unsatisfactory
	Human	PBMC	PMA	Cytofix	Perm I, II, or III	Greater induction on monocytes than lymphocytes, Perms I & II better than Perm III
WB	Human	HeLa S3	Anisomycin			25-kDa band induced
	Human	HeLa S3	Anisomycin + non-phospho peptide or alkaline phosphatase			No blocking of 25-kDa band
	Human	HeLa S3	Anisomycin + phospho peptide or λ phosphatase			Blocking of 25-kDa band
	Human	Jurkat or PBMC	PMA			25-kDa band induced

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Recommended Assay Procedure:

Either BD Cytofix™ fixation buffer or BD™ Phosflow Fix Buffer I may be used for cell fixation. Any of the three BD™ Phosflow permeabilization buffers may be used.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554655	Fixation Buffer	100 ml	(none)
557870	Fix Buffer I	250 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558052	Perm Buffer II	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Alexa Fluor® 647 fluorochrome emission is collected at the same instrument settings as for allophycocyanin (APC).
4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
5. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
6. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
7. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
8. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.

References

Culjkovic B, Topisirovic I, Borden KLB. Controlling gene expression through RNA regulons. The role of the eukaryotic translation initiation factor eIF4E. *Cell Cycle*. 2007; 6(1):65-69. (Biology)

Grolleau A, Kaplan MJ, Hanash SM, Beretta L, Richardson B. Impaired translational response and increased protein kinase PKR expression in T cells from lupus patients. *J Clin Invest*. 2000; 106(12):1561-1568. (Biology)

Mendez R, Myers Jr MG, White MF, Rhoads RE. Stimulation of protein synthesis, eukaryotic translation initiation factor 4E phosphorylation, and PHAS-I phosphorylation by insulin requires insulin receptor substrate 1 and phosphatidylinositol 3-kinase. *Mol Cell Biol*. 1996; 16(6):2857-2864. (Biology)

Waskiewicz AJ, Flynn A, Proud CG, Cooper JA. Mitogen-activated protein kinases activate the serine/threonine kinases Mnk1 and Mnk2. *EMBO J*. 1997; 16(8):1909-1920. (Biology)

Wendel H-G, Silva RLA, Malina A, et al. Dissecting eIF4E action in tumorigenesis. *Genes Dev*. 2007; 21:3232-3237. (Biology)