

## Technical Data Sheet

## PE Mouse anti-Human IRAK4


## Product Information

|                  |                                                                   |
|------------------|-------------------------------------------------------------------|
| Material Number: | 560303                                                            |
| Alternate Name:  | IRAK-4, REN64, Q69FE3                                             |
| Size:            | 50 tests                                                          |
| Vol. per Test:   | 20 µl                                                             |
| Clone:           | L29-525                                                           |
| Immunogen:       | Human IRAK4 Recombinant Protein                                   |
| Isotype:         | Mouse (BALB/c) IgG1, κ                                            |
| Reactivity:      | QC Testing: Human                                                 |
| Storage Buffer:  | Aqueous buffered solution containing BSA and ≤0.09% sodium azide. |


## Description

Interleukin 1 (IL-1) is a proinflammatory cytokine that mediates the host inflammatory response. IL-1 exerts its biological effects through its type I receptor (IL-1RI), which is found on most cell types. When IL-1 binds to the IL-1RI, NF-κB (transcription factor) is activated and translocates to the nucleus where it induces gene expression. IL-1-receptor-associated kinase (IRAK) has been reported to co-immunoprecipitate with IL-1RI. The IRAK family is made up of four identified Pelle homologs that regulate the activation of NF-κB and mitogen-activated protein (MAP) kinase pathways. IRAK4 is a unique member of the IRAK family because it is the closest mammalian homolog to Pelle. Also, when overexpressed, its kinase activity allows it to activate both the NF-κB and mitogen-activated protein (MAP) kinase pathways, and block the IL-1-induced activation and modification of IRAK1. IRAK4 is also able to phosphorylate IRAK1, which suggests that IRAK4 affects the early signal transduction of Toll/IL-1 receptors before the involvement of IRAK1.

The L29-525 monoclonal antibody recognizes human IRAK4, regardless of phosphorylation status. The specificity of this antibody conjugate for flow cytometric analysis was validated by confirming that RNA-mediated interference (RNAi) of the specific protein reduced the staining of the cells (see figure). Furthermore, the capacity of the RNAi to down-regulate the expression of the relevant protein was confirmed by western blot analysis.



**Analysis of IRAK4 in human peripheral blood lymphocytes.** Human peripheral blood mononuclear cells (PBMC) were lysed and fixed with 1X BD™ Phosflow Lyse/Fix Buffer (Cat. No. 558049) for 10 minutes at 37°C, then permeabilized (BD™ Phosflow Perm Buffer III, Cat. No. 558050) on ice for 30 minutes, and then stained with either PE Mouse IgG1, κ Isotype Control (Cat. No. 559320, shaded histogram) or PE Mouse anti-Human IRAK4 (open histogram). For data analysis, lymphocytes were selected by scatter profile. Flow cytometry was performed on a BD™ FACSCalibur flow cytometry system.



**Analysis of IRAK4 in human epithelioid carcinoma.** HeLa S3 cells (ATCC CCL 2.2) were either transfected with IRAK4 RNAi (open histogram) or untreated (shaded histogram). The cells were fixed (BD Cytofix™ buffer, Cat. No. 554655) for 10 minutes at 37°C, then permeabilized (BD™ Phosflow Perm Buffer III, Cat. No. 558050) on ice for 30 minutes, and then stained with PE Mouse anti-Human IRAK4. Flow cytometry was performed on a BD FACSCalibur™ flow cytometry system.

## BD Biosciences

bdbiosciences.com

|               |              |               |              |              |                         |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| United States | Canada       | Europe        | Japan        | Asia Pacific | Latin America/Caribbean |
| 877.232.8995  | 888.268.5430 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 0800.771.7157           |

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD




## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.



**Confirmation of the specificity of mAb L29-525 by western blot analysis (left panel).** Lysates of human PBMC were probed with unconjugated antibody at a concentration of 0.1 µg/ml. IRAK4 is identified as a band of about 52 kDa.

**Western blot validation of IRAK4 by RNAi (right panel).** Lysates from HeLa S3 cells (lane 1) and IRAK4 RNAi-transfected HeLa S3 cells (lane 2) were probed with unconjugated mAb L29-525 at 0.015 µg/ml. Down-regulation of IRAK4 expression is evident in the RNAi-transfected cells.

## Application Notes

### Application

|                                         |                  |
|-----------------------------------------|------------------|
| Intracellular staining (flow cytometry) | Routinely Tested |
|-----------------------------------------|------------------|

### Recommended Assay Procedure:

This antibody conjugate is suitable for intracellular staining of human peripheral blood mononuclear cells and cell lines (using BD™ Phosflow Lyse/Fix Buffer or BD Cytotfix™ Fixation Buffer), but not human whole blood. BD™ Phosflow Perm Buffer II or Perm Buffer III may be used.

### Suggested Companion Products

| Catalog Number | Name                             | Size      | Clone   |
|----------------|----------------------------------|-----------|---------|
| 558049         | Lyse/Fix Buffer 5X               | 250 ml    | (none)  |
| 554655         | Fixation Buffer                  | 100 ml    | (none)  |
| 558052         | Perm Buffer II                   | 125 ml    | (none)  |
| 558050         | Perm Buffer III                  | 125 ml    | (none)  |
| 559320         | PE Mouse IgG1, κ Isotype Control | 100 tests | MOPC-21 |

### Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use  $1 \times 10^6$  cells in a 100-µl experimental sample (a test).
2. Please refer to [www.bdbiosciences.com/pharminingen/protocols](http://www.bdbiosciences.com/pharminingen/protocols) for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

### References

- Hatao F, Muroi M, Hiki N, Ogawa T, Mimura Y, Kaminishi M, Tanamoto K. Prolonged Toll-like receptor stimulation leads to down-regulation of IRAK-4 protein. *J Leukoc Biol.* 2004; 76(4):904-908. (Biology)
- Li S, Strelow A, Fontana EJ, Wesche H. IRAK-4: a novel member of the IRAK family with the properties of an IRAK-kinase. *Proc Natl Acad Sci U S A.* 2002; 99(8):5567-5572. (Biology)
- Lye E, Mirtsos C, Suzuki N, Suzuki S, Yeh WC. The role of interleukin 1 receptor-associated kinase-4 (IRAK-4) kinase activity in IRAK-4-mediated signaling. *J Biol Chem.* 2004; 279(39):40653-40658. (Biology)
- Suzuki N, Saito T. IRAK-4 – a shared NF-κB activator in innate and acquired immunity. *Trends Immunol.* 2006; 27(12):566-572. (Biology)