

Technical Data Sheet

PE Mouse anti-MEK2

Product Information


Material Number:	560388
Alternate Name:	MAP Kinase Kinase 2; ERK Kinase
Size:	50 tests
Vol. per Test:	20 µl
Clone:	96/MEK2
Immunogen:	Rat MEK2 aa. 1-110 Recombinant Protein
Isotype:	Mouse IgG2a
Reactivity:	QC testing: Human Tested in development by western blot using purified antibody: chicken, dog, frog, mouse, rat
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

MEK (Map/Erk Kinase) 1 and 2 are serine/threonine kinases, also known as MAP kinase kinases (MAP2K1 and 2, MAPKK1 and 2, or MKK1 and 2). They activate the MAP (Mitogen-Activated Protein) kinases, also known as ERKs (Extracellular signal Regulated Kinases), which are critical kinases in multiple signal transduction pathways that regulate cell growth and differentiation. Activation of MEK 1 and 2 is dependent upon phosphorylation of serines 218 and/or 222 by activated MAP kinase kinases (MAP3Ks), such as the Raf isoforms. Hormones, growth and differentiating factors, or tumor promoters induce Raf activation via activation of Ras proteins. MEK2 is seven amino acids larger and shares 81% identity with MEK1. In cultured cells, MEK2 is activated by serum. In vitro, v-Raf phosphorylates and activates MEK2. It is thought that all of these activated protein kinases are downstream of the Ras signal transduction pathway and represent an integral part of the Ras mitogenic signal.

The 96/MEK2 monoclonal antibody recognizes MEK2, regardless of phosphorylation status.

The specificity of this antibody conjugate for flow cytometric analysis was validated by confirming that RNA-mediated interference (RNAi) of the specific protein reduced the staining of the cells (see figure).



LEFT: Analysis of MEK2 in human peripheral blood lymphocytes. Human peripheral blood mononuclear cells (PBMC) were lysed and fixed with BD Cytofix™ Fixation Buffer (Cat. No. 554655) for 10 minutes at 37°C, permeabilized (BD Phosflow™ Perm Buffer III, Cat. No. 558050) on ice for 30 minutes, and then stained with either PE Mouse IgG2a, κ Isotype Control (Cat. No. 558595, shaded histogram) or PE Mouse anti-MEK2 (open histogram). For data analysis, lymphocytes were selected by scatter profile. Flow cytometry was performed on a BD FACSArray™ flow cytometry system.

RIGHT: Analysis of MEK2 in human epithelioid carcinoma. HeLa S3 cells (ATCC CCL 2.2) were either transfected with MEK2 RNAi (shaded histogram) or untreated (open histogram). The cells were fixed (BD Cytofix™ buffer, Cat. No. 554655) for 10 minutes at 37°C, permeabilized (BD Phosflow™ Perm Buffer III, Cat. No. 558050) on ice for 30 minutes, and then stained with PE Mouse anti-MEK2. Flow cytometry was performed on a BD FACSArray™ flow cytometry system.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Application Notes

Application

Intracellular staining (flow cytometry)

Routinely Tested

Recommended Assay Procedure:

This antibody conjugate is suitable for intracellular staining of human cell lines and peripheral blood mononuclear cells using BD Cytotfix™ Fixation Buffer. Any of the three BD Phosflow™ permeabilization buffers may be used; Perm Buffers II and III are preferred. We do not recommend using it for staining of whole blood using BD Phosflow™ Lyse/Fix Buffer.

Suggested Companion Products

Catalog Number	Name	Size	Clone
610235	Purified Mouse Anti-MEK2	50 µg	96/MEK2
610236	Purified Mouse Anti-MEK2	150 µg	96/MEK2
558595	PE Mouse IgG2a, κ Isotype Control	50 tests	MOPC-173
554655	Fixation Buffer	100 ml	(none)
557885	Perm/Wash Buffer I	125 ml	(none)
558052	Perm Buffer II	125 ml	(none)
558050	Perm Buffer III	125 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100-µl experimental sample (a test).
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Crews CM, Alessandrini A, Erikson RL. The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science*. 1992; 258(5081):478-480. (Biology)

Downey GP, Butler JR, Tapper H, et al. Importance of MEK in neutrophil microbicidal responsiveness. *J Immunol*. 1998; 160(1):434-443. (Biology)

Hattori S, Fukuda M, Yamashita T, Nakamura S, Gotoh Y, Nishida E. Activation of mitogen-activated protein kinase and its activator by ras in intact cells and in a cell-free system. *J Biol Chem*. 1992; 267(28):20346-20351. (Biology)

Tworowski KA, Salghetti SE, Tansey WP. Stable and unstable pools of Myc protein exist in human cells. *Oncogene*. 2002; 21(55):8515-8520. (Biology)

Wu J, Harrison JK, Dent P, Lynch KR, Weber MJ, Sturgill TW. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK2. *Mol Cell Biol*. 1993; 13(8):4539-4548. (Biology)

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 800.979.9408 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

