Technical Data Sheet

FITC Mouse Anti-Human CD34

Product Information

Material Number:

gp105-120; My10; Hematopoietic progenitor cell antigen CD34 Alternate Name:


Size: 20 µl Vol. per Test: 581 Clone:

Mouse IgG1, κ Isotype: Reactivity: QC Testing: Human V MA27, VI E004 Workshop:

Aqueous buffered solution containing BSA and ≤0.09% sodium azide. Storage Buffer:

Description

The 581 monoclonal antibody specifically binds to CD34, a single-chain 105-120 kDa heavily O-glycosylated transmembrane glycoprotein expressed on a hemotopoietic progenitor cells, vascular endothelium and some tissue fibroblasts. The intracellular chain of the CD34 antigen is a target for phosphorylation by activated protein kinase C suggesting CD34 may play a role in signal transduction. CD34 may also play a role in adhesion of specific antigens to endothelium. Clone 581 reacts with the class III CD34 epitope, it is resistant to neuraminidase, chymopapain and glycoprotease. The 581 antibody blocks reactivity of another anti-CD34 monoclonal antibody, 8G12.



Profile of peripheral blood mononuclear cells analyzed by flow cytometry.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Application Notes

Application

Flow cytometry Routinely Tested

Suggested Companion Products

Catalog Number Clone 555748 FITC Mouse IgG1, κ Isotype Control 100 tests MOPC-21

Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 × 10⁶ cells in a 100-μl experimental
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 888.268.5430 32.53.720.550 0120.8555.90 65.6861.0633 0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



- 3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Egeland T, Tjonnfjord G, Steen R, Gaudernack G, Thorsby E. Positive selection of bone marrow-derived CD34 positive cells for possible stem cell transplantation. *Transplant Proc.* 1993; 25(1):1261-1263. (Biology)

Knapp W, Dorken B, Rieber EP, et al, ed. *Leucocyte Typing IV*. New York: Oxford University Press; 1989:1-1208. (Biology)

Schlossman SF, Bournsell L, Gilks W, et al, ed. Leukocyte Typing V: White Cell Differentiation Antigens. New York: Oxford University Press; 1995. (Biology)

560942 Rev. 1 Page 2 of 2