

Technical Data Sheet


V450 Mouse Anti-Human MIP-1 β **Product Information**

Material Number:	561282
Alternate Name:	Macrophage inflammatory protein 1-beta; CCL4; C-C motif chemokine 4; LAG-1
Size:	50 tests
Vol. per Test:	5 μ l
Clone:	D21-1351
Immunogen:	Recombinant Human MIP-1 β
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing protein stabilizer and \leq 0.09% sodium azide.

Description

The D21-1351 monoclonal antibody specifically binds to the human CC chemokine, MIP-1 β (macrophage inflammatory protein-1 β). Human MIP-1 β shares approximately 75% homology with mouse MIP-1 β at the amino acid level. Expression of MIP-1 β in human peripheral blood cells is induced by proinflammatory and mitogenic stimuli. MIP-1 β is a chemoattractant for monocytes and lymphocytes. Human MIP-1 β binds to receptors, CCR5 and CCR8. The human MIP-1 β gene has been mapped to chromosome 17q11. The immunogen used to generate D21-1351 hybridoma was recombinant human MIP-1 β .

The antibody is conjugated to BD Horizon™ V450, which has been developed for use in multicolor flow cytometry experiments and is available exclusively from BD Biosciences. It is excited by the Violet laser Ex max of 406 nm and has an Em Max at 450 nm. Conjugates with BD Horizon™ V450 can be used in place of Pacific Blue™ conjugates.



Flow cytometric analysis of MIP-1 β expressed in human peripheral blood mononuclear cells (PBMC). Human PBMC were either unstimulated (Left Panel) or stimulated (Right Panel) with 20 ng/mL Recombinant Human IFN- γ (Cat. No. 554616) for one hour followed by overnight incubation with 1 μ g/mL LPS (Sigma-Aldrich, Cat. No. L-8272) in the presence of 2 μ M BD GolgiStop™ Protein Transport Inhibitor (Cat. No. 554724). The PBMC were harvested, fixed with BD Cytotfix™ Fixation Buffer (Cat. No. 554655) and permeabilized with BD Perm/Wash™ Buffer (Cat. No. 554723). The cells were then stained with either a BD Horizon™ V450 Mouse IgG1, κ Isotype Control (Cat. No. 560373; dashed line histogram) or with the BD Horizon™ V450 Mouse Anti-Human MIP-1 β antibody (Cat. No. 561282; solid line histogram). The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable monocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ V450 under optimum conditions, and unreacted BD Horizon™ V450 was removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
560373	V450 Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21
554616	Recombinant Human IFN-γ	25 µg	(none)
554724	Protein Transport Inhibitor (Containing Monensin)	0.7 ml	(none)
554655	Fixation Buffer	100 ml	(none)
554723	Perm/Wash Buffer	100 ml	(none)
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100-µl experimental sample (a test).
2. An isotype control should be used at the same concentration as the antibody of interest.
3. Please refer to wwwbdbiosciences.com/pharmingen/protocols for technical protocols.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at wwwbdbiosciences.com/colors.
6. BD Horizon™ V450 has a maximum absorption of 406 nm and maximum emission of 450 nm. Before staining with this reagent, please confirm that your flow cytometer is capable of exciting the fluorochrome and discriminating the resulting fluorescence.
7. Pacific Blue™ is a trademark of Molecular Probes, Inc., Eugene, OR.

References

- Bernardini G, Hedrick J, Sozzani S. Identification of the CC chemokines TARC and macrophage inflammatory protein-1 beta as novel functional ligands for the CCR8 receptor. *J Immunol.* 1998; 28(2):582-588. (Biology)
- Combadiere C, Ahuja SK, Tiffany HL, Murphy PM. Cloning and functional expression of CC CKR5, a human monocyte CC chemokine receptor selective for MIP-1(alpha), MIP-1(beta), and RANTES. *J Leukoc Biol.* 1996; 60(1):147-152. (Biology)
- Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods.* 1995; 188(1):117-128. (Methodology: Flow cytometry, IC/FCM Block)
- Raport CJ, Gosling J, Schweickart VL, Gray PW, Charo IF. Molecular cloning and functional characterization of a novel human CC chemokine receptor (CCR5) for RANTES, MIP-1beta, and MIP-1alpha. *J Biol Chem.* 1996; 271(29):17161-17166. (Biology)