BD™ Cytometric Bead Array (CBA)



Technical Data Sheet

Mouse IL-5 Enhanced Sensitivity Flex Set

Product Information

- Material Number: Size: Bead Position: Assay Range: Reactivity:
- 562234 100 Tests A6 274-200,000 fg/mL QC Testing: Mouse

Component Description: Component Mat. No: Component Storage Buffer:

Component Description: Component Mat. No: Component Storage Buffer:

Component Description: Component Mat. No: Component Storage Buffer: 51-9003524 Lyophilized in an aqueous buffered solution containing BSA and ProClin[™] 150. Mouse IL-5 Enhanced Sensitivity Capture Bead A6

Mouse IL-5 Standard

51-9007636 Aqueous buffered solution containing fetal bovine serum and ≤0.09% sodium azide.

Mouse IL-5 Detection Reagent (Part A) 51-9007637 Aqueous buffered solution containing BSA and ProClin[™] 150 as preservative.




Figure 1. Example of BD CBA Mouse IL-5 Enhanced Sensitivity Flex Set standard curve. Data was acquired using a BD FACSArray™ Bioanalyzer and analyzed using FCAP Array Software (Cat. No. 641488).

Description

The BDTM CBA Mouse IL-5 Enhanced Sensitivity Flex Set is a bead-based immunoassay capable of measuring mouse Interleukin-5 (IL-5) in serum and cell culture supernatant samples. The BD CBA Enhanced Sensitivity Flex Sets are capable of measuring soluble analytes present in very low concentrations. For more information on bead-based immunoassays, refer to the product insert for the BD CBA Mouse Enhanced Sensitivity Master Buffer Kit (Cat. No. 562246 or 562248).

This BD CBA Enhanced Sensitivity Flex Set contains one vial each of Enhanced Sensitivity Capture Bead and Detection Reagent (Part A) and two vials of Standard. The Enhanced Sensitivity Detection Reagent (Part B) is provided in the BD CBA Mouse Enhanced Sensitivity Master Buffer Kit. The Enhanced Sensitivity Capture Bead and Detection Reagent (Part A) components of this flex set have been formulated to a 20x concentration to ensure product performance when multiplexed. The Standard component is lyophilized and when reconstituted according to the instructions in the BD CBA Mouse Enhanced Sensitivity Master Buffer kit, the top standard point for this assay is 200,000 fg/ml. Discard unused reconstituted standard, do not store or reuse. Store lyophilized standard and other components at 4°C.

Application Notes

Recommended Assay Procedure: The BD CBA Mouse IL-5 Enhanced Sensitivity Flex Set must be used in conjunction with a BD CBA Mouse Enhanced Sensitivity Master Buffer Kit (Cat. No. 562246, 100 tests, or 562248, 500 tests), a flow cytometer, and the FCAP Array™ Software (Cat. No. 641488). Detailed instructions on the use of this product can be found in the manual for the BD CBA Mouse Enhanced Sensitivity Master Buffer Kit. When following the directions in the Master Buffer Kit, the standard range for the BD CBA Mouse IL-5 Enhanced Sensitivity Flex Set will be 274 to 200,000 fg/ml. An example standard curve is shown in Figure 1.

The BD CBA Human IL-5 Enhanced Sensitivity Flex Set should only be used in the same assay well with other BD CBA Mouse Enhanced Sensitivity Flex Set reagents. We do not recommend plexing Enhanced Sensitivity Flex Set assays along with assays from any other Flex Set

BD Biosciences

bdbiosciences.c					
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995
For country cor	ntact informatio	on, visit bdbiosci e	ences.com/conta	ct	
of any patents. BE use of our produc product or as a co written authoriza For Research Use (Delissciences will not ts. Purchase does not mponent of anoth- tion of Becton, Dick Dnly. Not for use in	ot be held responsik ot include or carry a er product. Any use kinson and Compan diagnostic or thera	ole for patent infrin any right to resell or of this product oth y is stictly prohibite peutic procedures.	gement or other vio r transfer this produce er than the permitte d. Not for resale.	a the above product in violation lations that may occur with the ct either as a stand-alone ad use without the express n and Company. © 2014 BD



system (soluble protein, cell signaling, etc.). For an updated assay compatibility chart for the BD CBA Mouse Enhanced Sensitivity Flex Sets, please refer to the BD CBA Flex Set System homepage at http://www.bdbiosciences.com/cbasetup.

Performance

Limit of Detection: The theoretical limit of detection is 12.5 fg/ml and was determined by evaluating the estimated result of the average MFI of the negative control (0 fg/ml, n=30) + 2 standard deviations.

Specificity		Inter-Assay Reproducibility			Intra-Assay Reproducibility		
		Mean (fg/ml)	Standard Deviation	%CV	Mean (fg/ml)	Standard Deviation	%CV
	Sample 1	2,057.9	210.7	10%	1,742.1	96.0	6%
Mouse IL-5	Sample 2	6,152.4	813.9	13%	4,812.2	239.8	5%
	Sample 3	19,625.8	2,835.7	14%	15,282.3	644.1	4%

Reproducibility: The intra-assay and inter-assay reproducibility were determined for the BD CBA Mouse IL-5 Enhanced Sensitivity Flex Set by evaluating ten replicates of three different sample levels (intra-assay) and three replicates of three different sample levels from four separate experiments (inter-assay) respectively.

		ulture natant	Serum		
Specificity	Average % Recovery	Range	Average % Recovery	Range	
Mouse IL-5	73%	68 - 80%	65%	62 - 67%	

Recovery: Cell culture supernatant and pooled mouse serum were spiked with three different levels of protein. The spiked samples were assayed and the results were compared with expected values. Serum samples were diluted 1:3 before the protein was spiked into each. Serum used was commercially available pooled mouse serum.

		ulture natant	Serum		
Sample Dilution	Detected (fg/ml)	% of Expected	Detected (fg/ml)	% of Expected	
Spiked sample	16,898.7	100%	15,607.6	100%	
1:4	5,259.5	93%	5,264.3	101%	
1:8	1,821.0	97%	1,847.9	107%	

Linearity: Cell culture supernatant and 1:3 diluted pooled mouse serum were spiked with protein and serially diluted. The diluted samples were assayed and the results were compared with the original spiked sample.

Product Notices

- 1. ProClin is a trademark of Rohm and Haas Company.
- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Warning: CBA lyophilized standard contains 0.02% (w/w) and Detection Reagent (Part A) contains 0.002% (w/w) of a CMIT/MIT mixture (3:1), which is a mixture of: 5-chloro-2-methyl-4-isothiazolin-3-one [EC No 247-500-7] and 2-methyl-4-isothiazolin-3-one [EC No 220-239-6] (3:1).

Hazard statement: May cause an allergic skin reaction.

Precautionary statements: Wear protective gloves/eye protection. Wear protective clothing. Avoid breathing mist/vapours/spray. If skin irritation or rash occurs: Get medical advice/attention. IF ON SKIN: Wash with plenty of water. Dispose of contents/container in accordance with local/regional/national/international regulations.

bdbiosciences.com							
United States 877.232.8995	Canada 800.268.5430	Europe 32.2.400.98.95	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbear 55.11.5185.9995		
For country cor	ntact informatio	on, visit bdbiosci e	ences.com/conta	ct			
of any patents. BL use of our produc product or as a co	D Biosciences will ne ts. Purchase does n mponent of anoth	ot be held responsit ot include or carry a	ble for patent infrin any right to resell of of this product oth	gement or other vio r transfer this producer than the permitte	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express		

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

