Technical Data Sheet

BV421 Hamster Anti-Mouse CD95

Product Information

Material Number:	562633		
Alternate Name:	Apo-1; Apt1; Fas; FASLG receptor; lpr; TNFR6; Tnfrsf6; TNR6		
Size:	50 µg		
Concentration:	0.2 mg/ml		
Clone:	Jo2		
Immunogen:	WR19L mouse lymphoma cells transformed with recombinant mouse Fas		
Isotype:	Armenian Hamster IgG2, $\lambda 2$		
Reactivity:	QC Testing: Mouse		
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.		

Description

Fas antigen, CD95, is a 45 kDa cell-surface protein which can mediate apoptosis. It belongs to the TNF (tumor necrosis factor)/NGF receptor family. Expression of Fas has been described in the thymus, liver, heart, lung and ovary. Fas plays an important role in the apoptotic process that takes place during development. Monoclonal antibodies recognizing Fas such as Jo2 have cytolytic activity on cells expressing Fas. The cell death stimulated by Fas antibodies is characteristic of apoptosis and suggests that the lethal effects are a result of interaction of antibody with a functional Fas antigen as opposed to complement-mediated lysis.

The Jo2 antibody recognizes mouse Fas. The Jo2 antibody shows cytolytic activity against cell lines expressing mouse Fas by inducing apoptosis. Intraperitoneal injections of Jo2 mAb have been shown to kill mice and induce apoptotic hepatocyte death. Jo2 mAb has been reported to immunoprecipitate mouse Fas as a 45 kDa band from W4 cells. W4 cells are WR19L mouse lymphoma cells transformed with mouse Fas. The difference between the observed MW of Fas and that deduced from its amino acid sequence (Mr 34,971) may be due to glycosylation.

The antibody was conjugated to BD HorizonTM BV421 which is part of the BD HorizonTM Brilliant VioletTM family of dyes. With an Ex Max of 407-nm and Em Max at 421-nm, BD HorizonTM BV421 can be excited by the violet laser and detected in the standard Pacific BlueTM filter set (eg, 450/50-nm filter). BD Horizon™ BV421 conjugates are very bright, often exhibiting a 10 fold improvement in brightness compared to Pacific Blue[™] conjugates.



Flow cytometric analysis of CD95 expression on mouse thymocytes. Thymocytes from a BALB/c mouse were stained with either a BD Horizon™ BV421 Hamster IgG2, λ1 isotype control (Cat. No. 562629; dashed line histogram) or with the BD Horizon™ BV421 Hamster Anti-Mouse CD95 antibody (Cat. No. 562633; solid line histogram). The fluorescence histograms were derived from events with the forward and side light-scatter characteristics of viable thymocytes. Flow cytometry was performed using a BD™ LSR II Flow Cytometer System.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD HorizonTM BV421 under optimum conditions, and unconjugated antibody and free BD HorizonTM BV421 were removed.

BD Biosciences

bdbiosciences.com									
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean				
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995				
For country contact information, visit bdbiosciences.com/contact									
Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation									
of any patents. Bl	D Biosciences will n	ot be help responsi	ble for patent infrin	ngement or other vio	lations that may occur with the				
use of our produc	ts. Purchase does n	ot include or carry	any right to resell o	r transfer this produ	ct either as a stand-alone				
product or as a component of another product. Any use of this product other than the permitted use without the every									

written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Application Notes

A	Аррисатоя								
	Flow cytometry	Routinely Tested							
S	Suggested Companion Products								

Catalog Number	Name	Size	Clone
562629	BV421 Hamster IgG2, Lambda Isotype Control	50 μg	Ha4/8
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

Since applications vary, each investigator should titrate the reagent to obtain optimal results. 1.

- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. An isotype control should be used at the same concentration as the antibody of interest.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at 6. www.bdbiosciences.com/colors.
- Pacific Blue[™] is a trademark of Molecular Probes, Inc., Eugene, OR. 7.
- Brilliant Violet[™] 421 is a trademark of Sirigen. 8.
- Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian 9. hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/documents/hamster_chart_11x17.pdf.

References

Hiromatsu K, Aoki Y, Makino M, et al. Increased Fas antigen expression in murine retrovirus-induced immunodeficiency syndrome, MAIDS. Eur J Immunol. 1994; 24(10):2446-2451. (Clone-specific: Cytotoxicity, Flow cytometry)

Kagi D, Vignaux F, Ledermann B, et al. Fas and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. Science. 1994; 265(5171):528-530. (Clone-specific: Flow cytometry, Functional assay)

Ogasawara J, Suda T, Nagata S. Selective apoptosis of CD4+CD8+ thymocytes by the anti-Fas antibody. J Exp Med. 1995; 181(2):485-491. (Clone-specific: Flow cvtometry, Functional assay)

Ogasawara J, Watanabe-Fukunaga R, Adachi M, et al. Lethal effect of the anti-Fas antibody in mice. Nature. 1993; 364(6440):806-809. (Immunogen: Flow cytometry, Immunoprecipitation)

Yang Y, Mercep M, Ware CF, Ashwell JD. Fas and activation-induced Fas ligand mediate apoptosis of T cell hybridomas: inhibition of Fas ligand expression by retinoic acid and glucocorticoids. J Exp Med. 1995; 181(5):1673-1682. (Clone-specific: Flow cytometry)

BD Biosciences

bdbiosciences.com United States Canada 877.232.8995

Europe Japan 800.979.9408 32.53.720.550 0120.8555.90 For country contact information, visit bdbiosciences.com/contact

Latin America/Caribbean 65.6861.0633 55.11.5185.9995



Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be help responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Asia Pacific