

Technical Data Sheet

Purified Rat anti-Human Lgr5 (CRL Region)


Product Information

Material Number:	562733
Alternate Name:	GPR49, GPR67, HG38
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	1D9A5 (also known as 1D9)
Immunogen:	Human LGR5 DNA
Isotype:	Rat IgG2b, λ
Reactivity:	QC Testing: Human Not Reactive: Mouse
Target MW:	~110 kDa
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.


Description

Lgr5 (leucine-rich-repeat-containing G-protein-coupled receptor 5) is a seven transmembrane-domain receptor that is a target gene for Wnt and marks stem cells in the small intestine and colon, stomach, and hair follicle. Lgr5 was initially identified as a potential stem cell marker due to restricted expression of Lgr5 in the intestinal crypt and labeling of rapidly cycling cells of the colon and intestine. Using both lineage tracing and organoid culture experiments, Lgr5 positive cells are capable of generating all types of the small intestine epithelium hence indicating that Lgr5 marks stem cells of the small intestine and colon. R-spondin growth factors, which are secreted agonists of the Wnt pathway, bind Lgr5. The binding of R-spondins to Lgr5 leads to recruitment of the Frizzled/LRP Wnt receptor complex, which binds to Wnt ligands and leads to downstream Wnt signaling. Lgr5 is up-regulated in colon and ovarian cancers and has been implicated in promotion of tumor growth and metastasis.

The 1D9A5 monoclonal antibody recognizes an epitope in the Cysteine-rich linker (CRL) region of Human Lgr5.



Western blot analysis of human Lgr5. Lysate from colorectal adenocarcinoma cells LS 174T transfected with human LGR5 (Cells from Dr. Hans Clevers, Hubrecht Institute) was probed with Purified Rat anti-Human Lgr5 (CRL Region) monoclonal antibody (Cat. No. 562733) at titrations of 1.0 (lane 1), 0.5 (lane 2), and 0.25 $\mu\text{g/ml}$ (lane 3). Lgr5 is identified as a strong band of ~110 kDa. The protein was detected using HRP Goat Anti-Rat Ig (Cat. No. 554017) and a chemiluminescent detection system.



Immunofluorescent staining of human Lgr5. Colorectal adenocarcinoma cells LS 174T transfected with human LGR5 (Cells from Dr. Hans Clevers, Hubrecht Institute) were stained with Purified Rat anti-Human Lgr5 (CRL Region) monoclonal antibody (Cat. No. 562733) (pseudo colored green) at 5 $\mu\text{g/mL}$. The second-step reagent was Alexa Fluor® 488 goat anti-Rat IgG (Life Technologies). The images were captured on a BD Pathway™ 435 Cell Analyzer and merged using BD Attovision™ Software.

Preparation and Storage

Store undiluted at 4°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

Application Notes

Application

Western blot	Routinely Tested
Flow cytometry	Tested During Development
Bioimaging	Tested During Development
Immunofluorescence	Tested During Development
Immunohistochemistry-frozen	Not Recommended
Immunohistochemistry-formalin (antigen retrieval required)	Not Recommended

Suggested Companion Products

Catalog Number	Name	Size	Clone
353219	BD Falcon™ 96-well Imaging Plate		(none)
554017	HRP Goat Anti-Rat Ig	1.0 ml	Polyclonal
550767	PE Goat Anti-Rat Ig	0.2 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE (No Azide/Low Endotoxin) antibody format, if available, for in vitro and in vivo use.
5. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.

References

Barker N, Huch M, Kujala P, et al. Lgr5(+ve) stem cells drive self-renewal in the stomach and build long-lived gastric units in vitro. *Cell Stem Cell*. 2010; 6(1):25-36. (Biology)

Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, et al. Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature*. 2007; 449(7165):1003-1007. (Biology)

Carmon KS, Gong X, Lin Q, Thomas A, Liu Q. R-spondins function as ligands of the orphan receptors LGR4 and LGR5 to regulate Wnt/β-catenin signaling. *Proc Natl Acad Sci U S A*. 2011; 108(28):11452-11457. (Biology)

de Lau W, Barker N, Low TY, et al. Lgr5 homologues associate with Wnt receptors and mediate R-spondin signalling. *Nature*. 2011; 476(7360):293-297. (Clone-specific)

Jaks V, Barker N, Kasper M, et al. Lgr5 marks cycling, yet long-lived, hair follicle stem cells. *Nat Genet*. 2008; 40(11):1291-1299. (Biology)

Kemper K, Prasetyanti PR, de Lau W, Rodermond H, Clevers H, Medema JP. Monoclonal Antibodies Against Lgr5 Identify Human Colorectal Cancer Stem Cells. *Stem Cells*. 2012; . (Clone-specific: Flow cytometry, Immunofluorescence)

Merlos-Suárez A, Barriga FM, Jung P et al. The intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease relapse. *Cell Stem Cell*. 2011; 8(5):511-524. (Biology)

Yui S, Nakamura T, Sato T, et al. Functional engraftment of colon epithelium expanded in vitro from a single adult Lgr5(+) stem cell. *Nat Med*. 2012; 18(4):618-623. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

