

Technical Data Sheet

BV711 Rat Anti-Mouse CD326


Product Information

Material Number:	563134
Alternate Name:	Ep-CAM; EGP; EGP-2; Egp314; GA733-2; TROP1; Tacsd1; Tacstd1; Ly74; gp40
Size:	50 µg
Concentration:	0.2 mg/ml
Clone:	G8.8
Immunogen:	Glycoconjugates from BALB/c mouse-derived TE-71 medullary thymic epithelial cell line
Isotype:	Rat (SD) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The G8.8 antibody reacts with CD326/Ep-CAM (*Epithelial Cell Adhesion Molecule*), also known as gp40 in the mouse and by a variety of names (including GA733-2, CO17-1A, and EGP) in the human. In the mouse, Ep-CAM is a 40-42 kDa cell-surface type 1 transmembrane glycoprotein expressed on thymic epithelial cells, thymic dendritic cells, immature thymocytes, a small subset of peripheral T lymphocytes, intestinal epithelium, kidney-collecting tubule epithelium, keratinocytes, Langerhans cells and lymph node and splenic dendritic cells. Profiles of Ep-CAM expression on fetal thymocytes and on the CD4[-] CD8[-], CD4[+] CD8[+], CD4[-] CD8[+], and CD4[+] CD8[-] subsets of adult thymocytes have been published. In unrelated studies, mouse Ep-CAM mRNA was detected in tissues containing epithelial cells (kidney, stomach, intestine, lung, and thymus) and in plasma cells and plasmacytomas, but not in heart, muscle, liver, brain, spleen, B lymphomas, or pre-B lymphomas. Ep-CAM is a Ca²⁺ independent homophilic adhesion molecule that is proposed to play roles in the development and normal function of epithelial tissues and in the progression of carcinomas.

The antibody was conjugated to BD Horizon™ BV711 which is part of the BD Horizon™ Brilliant Violet™ family of dyes. This dye is a tandem fluorochrome of BD Horizon™ BV421 with an Ex Max of 405-nm and an acceptor dye with an Em Max at 711-nm. BD Horizon™ BV711 can be excited by the violet laser and detected in a filter used to detect Cy™5.5 / Alexa Fluor® 700-like dyes (eg, 712/20-nm filter). Due to the excitation and emission characteristics of the acceptor dye, there may be moderate spillover into the Alexa Fluor® 700 and PerCP-Cy™5.5 detectors. However, the spillover can be corrected through compensation as with any other dye combination.



Multicolor analysis of CD326 expression on mouse thymocytes and splenic T lymphocytes. BALB/c mouse thymocytes and splenic leucocytes were preincubated with Purified Rat Anti-Mouse CD16/CD32 antibody (Mouse BD Fc Block™) (Cat. No. 553141/553142) and stained with BD Horizon™ BV711 Rat IgG2a, κ Isotype Control (Cat. No. 563047) or BD Horizon™ BV711 Rat Anti-Mouse CD326 antibody (Cat. No. 563134).

Left Panel: Thymocytes were further stained with FITC Rat Anti-Mouse CD4 (553047/553046/561835) and PE Rat Anti-Mouse CD8a (553033/553032/561095) antibodies. The CD326 (solid line) and Ig Isotype Control (dashed line) fluorescence histograms were derived from CD4- and CD8-negative gated events with the forward and side light-scatter characteristics of viable thymocytes.

Middle and Right Panels: The splenic leucocytes were further stained with FITC Hamster Anti-Mouse CD3e (Cat. No. 553062/553061/561827) and PE Rat Anti-Mouse CD25 (Cat. No. 553866/561065) antibodies. Two-color flow cytometric dot plots show the correlated expression patterns of Ig Isotype control staining (Middle Plot) or CD326 (Right Plot) versus CD3e for CD25+ gated events with the forward and side light-scatter characteristics of viable splenic leucocytes. A small population of CD3+CD25+CD326+ cells was detected (Right Panel), whereas the CD25- T cells do not express detectable levels of CD326 (data not shown).

Flow cytometric analysis was performed using a BD™ LSR II Flow Cytometer System.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD Horizon™ BV711 under optimum conditions, and unconjugated antibody and free BD Horizon™ BV711 were removed.

Application Notes

Application

Flow cytometry

Routinely Tested

Suggested Companion Products

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 ml	(none)
563047	BV711 Rat IgG2a, κ Isotype Control	50 µg	R35-95
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2
553142	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.5 mg	2.4G2
555899	Lysing Buffer	100 ml	(none)
553062	FITC Hamster Anti-Mouse CD3e	0.5 mg	145-2C11
553061	FITC Hamster Anti-Mouse CD3e	0.1 mg	145-2C11
561827	FITC Hamster Anti-Mouse CD3e	25 µg	145-2C11
553866	PE Rat Anti-Mouse CD25	0.2 mg	PC61
561065	PE Rat Anti-Mouse CD25	25 µg	PC61
553047	FITC Rat Anti-Mouse CD4	0.5 mg	RM4-5
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
561835	FITC Rat Anti-Mouse CD4	25 µg	RM4-5
553033	PE Rat Anti-Mouse CD8a	0.2 mg	53-6.7
553032	PE Rat Anti-Mouse CD8a	0.1 mg	53-6.7
561095	PE Rat Anti-Mouse CD8a	25 µg	53-6.7

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
3. An isotype control should be used at the same concentration as the antibody of interest.
4. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
5. Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Eugene, OR.
6. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
7. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
8. Cy is a trademark of Amersham Biosciences Limited.
9. Brilliant Violet™ 711 is a trademark of Sirigen.

References

Basak S, Speicher D, Eck S, Wunner W. Colorectal carcinoma invasion inhibition by CO17-1A/GA733 antigen and its murine homologue. *J Natl Cancer Inst.* 1998; 90(9):691-697. (Biology)

Bergsagel PL, Victor-Kobrin C, Timblin CR, Trepel J, Kuehl WM. A murine cDNA encodes a pan-epithelial glycoprotein that is also expressed on plasma cells. *J Immunol.* 1992; 148(2):590-596. (Biology)

Birebent B, Somasundaram R, Purev E. Anti-idiotypic antibody and recombinant antigen vaccines in colorectal cancer patients. *Crit Rev Oncol Hematol.* 2001; 39(1-2):107-113. (Biology)

Borkowski TA, Nelson AJ, Farr AG, Udey MC. Expression of gp40, the murine homologue of human epithelial cell adhesion molecule (Ep-CAM), by murine dendritic cells. *Eur J Immunol.* 1996; 26(1):110-114. (Clone-specific: Immunohistochemistry)

Farr A, Nelson A, Truex J, Hosier S. Epithelial heterogeneity in the murine thymus: a cell surface glycoprotein expressed by subcapsular and medullary epithelium. *J Histochem Cytochem.* 1991; 39(5):345-353. (Immunogen: Electron microscopy, Immunohistochemistry, Immunoprecipitation)

Litvinov SV, Balzar M, Winter MJ. Epithelial cell adhesion molecule (Ep-CAM) modulates cell-cell interactions mediated by classic cadherins. *J Biol Chem.* 1997; 272(5):1337-1348. (Biology)

Nelson AJ, Dunn RJ, Peach R, Aruffo A, Farr AG. The murine homolog of human Ep-CAM, a homotypic adhesion molecule, is expressed by thymocytes and thymic epithelial cells. *Eur J Immunol.* 1996; 26(2):401-408. (Clone-specific: Electron microscopy, Immunohistochemistry, Immunoprecipitation)

Taguchi N, Hashimoto Y, Naiki M. Abnormal thymic expression of epithelial cell adhesion molecule (EP-CAM) in New Zealand Black (NZB) mice. *J Autoimmun.* 1999; 13(4):393-400. (Clone-specific: Electron microscopy)

Zutter MM. Gastrointestinal carcinoma antigen GA733: target for immunodestruction and potential modifier of invasiveness and chemoresponsiveness. *J Natl Cancer Inst.* 1998; 90(9):642-644. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

