Technical Data Sheet

BV605 Rat Anti-Mouse Vα2 TCR

Product Information


Material Number:	563286
Alternate Name:	T cell receptor alpha, variable region 2
Size:	50 µg
Concentration:	0.2 mg/ml
Clone:	B20.1
Immunogen:	Soluble a BTCR from mouse cytotoxic T-cell clone KB5-C20
Isotype:	Rat (LOU) IgG2a, λ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

The B20.1 monoclonal antibody specifically binds to most members of the Va2 T-cell Receptor (TCR) subfamily in mice having the a, b, and c haplotypes of the Tcrb gene complex. B20.1 antibody may crossreact with V δ 8 TCR, which shares >90% sequence homology with V α 2 TCR. Levels of B20.1+T cells appear to be influenced by V α haplotypes. Moreover, the frequencies of V α 2+ CD8+ and CD4+ T cells are influenced by H-2 haplotypes.

This antibody is conjugated to BD Horizon BV605 which is part of the BD Horizon Brilliant™ Violet family of dyes. With an Ex Max of 407-nm and Em Max of 602-nm, BD Horizon BV605 can be excited by a violet laser and detected with a standard 610/20-nm filter set. BD Horizon BV605 is a tandem fluorochrome of BD Horizon BV421 and an acceptor dye with an Em max at 605-nm. Due to the excitation of the acceptor dye by the green (532 nm) and yellow-green (561 nm) lasers, there will be significant spillover into the PE and BD Horizon PE-CF594 detectors off the green or yellow-green lasers. BD Horizon BV605 conjugates are very bright, often exhibiting brightness equivalent to PE conjugates and can be used as a third color off of the violet laser.

For optimal and reproducible results, BD Horizon Brilliant Stain Buffer should be used anytime two or more BD Horizon Brilliant dyes are used in the same experiment. Fluorescent dye interactions may cause staining artifacts which may affect data interpretation. The BD Horizon Brilliant Stain Buffer was designed to minimize these interactions. More information can be found in the Technical Data Sheet of the BD Horizon Brilliant Stain Buffer (Cat. No. 563794).



Two-color flow cytometric analysis of Va2 TCR expression on mouse lymph node cells. Lymph node cells from a BALB/c mouse were preincubated with Purified Rat Anti-Mouse CD16/CD32 antibody (Mouse BD Fc Block™) (Cat. No. 553141/553142). The cells were then stained with FITC Rat anti-Mouse CD4 (Cat. No. 553729/557307/561828) and FITC Rat anti-Mouse CD8b (Cat. No. 553040/561968) in addition to either BD Horizon™ BV605 Rat IgG2a, λ isotype control (Cat. No. 562998; Left Panel) or with BD Horizon™ BV605 Rat Anti-Mouse Vα2 TCR antibody (Cat. No. 563286; Right Panel). Two-color flow cytometric dot plots showing the correlated expression of Va2 TCR (or Ig Isotype control staining) versus CD4 and CD8b were derived from gated events with the forward and side light-scattering characteristics of viable lymph node cells. Flow cytometric analysis was performed using a BD™ LSR II Flow Cytometry System.

BD Biosciences

bablosciences.c	om					
United States 877.232.8995	Canada 800.268.5430	Europe 32.2.400.98.95	Japan	Asia Pacific 65.6861.0633	Latin America/Caribbean 55.11.5185.9995	
					55.11.5165.9995	
,		,	ences.com/contac			
of any patents. BD use of our product product or as a co	Biosciences will ne ts. Purchase does n mponent of anoth	ot be held responsik ot include or carry a er product. Any use	le for patent infring ny right to resell or t	ement or other viol transfer this produc r than the permitte	the above product in violation lations that may occur with the t either as a stand-alone d use without the express	

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with BD HorizonTM BV605 under optimum conditions, and unconjugated antibody and free BD HorizonTM

BV605 were removed. **Application Notes**

liaati

A	pplication
	Flow cytometry

9	Suggested Companion Products	
	euggeeteu eempunen rieuuete	

Catalog Number	Name	Size	Clone
554656	Stain Buffer (FBS)	500 mL	(none)
562998	BV605 Rat IgG2a, λ Isotype Control	50 µg	B39-4
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2
553142	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.5 mg	2.4G2
563794	Brilliant Stain Buffer	5 mL	(none)

Routinely Tested

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results. 1.
- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- An isotype control should be used at the same concentration as the antibody of interest. 3.
- 4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

5. Please observe the following precautions: Absorption of visible light can significantly alter the energy transfer occurring in any tandem fluorochrome conjugate; therefore, we recommend that special precautions be taken (such as wrapping vials, tubes, or racks in aluminum foil) to prevent exposure of conjugated reagents, including cells stained with those reagents, to room illumination.

- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 6. discarding to avoid accumulation of potentially explosive deposits in plumbing.
- For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at 7. www.bdbiosciences.com/colors.
- Although every effort is made to minimize the lot-to-lot variation in the efficiency of the fluorochrome energy transfer, differences in the 8 residual emission from BD Horizon[™] BV421 may be observed. Therefore, we recommend that individual compensation controls be performed for every BD Horizon[™] BV605 conjugate.
- CF[™] is a trademark of Biotium, Inc. 9.

References

Gregoire C, Rebai N, Schweisguth F, et al. Engineered secreted T-cell receptor alpha beta heterodimers. Proc Natl Acad Sci U S A. 1991; 88(18):8077-8081. (Immunogen: Flow cytometry, Immunoprecipitation, Radioimmunoassay)

Pircher H, Rebai N, Groettrup M, et al. Preferential positive selection of V alpha 2+ CD8+ T cells in mouse strains expressing both H-2k and T cell receptor V alpha a haplotypes: determination with a V alpha 2-specific monoclonal antibody. Eur J Immunol. 1992; 22(2):399-404. (Immunogen: Flow cytometry, Immunofluorescence. Immunoprecipitation)

Tomonari K, Fairchild S, Rosenwasser OA. Influence of viral superantigens on V beta- and V alpha-specific positive and negative selection. Immunol Rev. 1993; 131:131-168. (Methodology)

BD Biosciences

bdbiosciences.com United States

 Canada
 Europe
 Japan

 800.268.5430
 32.2.400.98.95
 0120.8555.90
Asia Pacific Latin America/Caribbean 877.232.8995 65.6861.0633 55.11.5185.999 For country contact information, visit bdbiosciences.com/contact

Conditions: The information insclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

